

RESEARCH ARTICLE

Open Access

VEGF is an important mediator of tumor angiogenesis in malignant lesions in a genetically engineered mouse model of lung adenocarcinoma

Bharat K Majeti[†], Joseph H Lee[†], Brett H Simmons[†] and Farbod Shojaei^{*}

Abstract

Background: VEGF is one of the key drivers of physiological or pathological angiogenesis hence several VEGF inhibitors are in different stages of clinical development. To further dissect the role of VEGF in different stages of tumor progression in lung tumors, we utilized $Kras^{G12D-LSL}$ GEMMs (genetically engineered mouse models).

Methods: Intranasal delivery of adenoviruses expressing cre recombinase in $Kras^{G12D-LSL}$ mice results in the expression of mutant *Kras* that leads to development of tumor lesions ranging from adenomatous hyperplasia to large adenoma and adenocarcinoma over time in lung. In the current study, we treated $Kras^{G12D-LSL}$ mice at 14 weeks post inhalation with three different angiogenic inhibitors including axitinib and PF-00337210 both of which are selective inhibitors of VEGFR and sunitinib which targets VEGFR, C-SF1-R, PDGFR and KIT.

Results: Pathology findings showed no significant difference in percentage of adenomatous hyperplastic lesions between the vehicle vs. any of the treatments suggesting that angiogenesis may not play a major role at early stages of tumorigenesis. However, each inhibitor suppressed percentage of benign adenoma lesions and almost fully inhibited growth of adenocarcinoma lesions in the recipients which was consistent with a reduction in tumor vasculature. Treatment with sunitinib which is a multi-targeted RTKI did not provide any advantage compared to selective VEGFR inhibitor further emphasizing role of VEGF in tumor angiogenesis in this model.

Conclusion: Overall, our studies indicate significance of VEGF and angiogenesis in a spontaneous model of lung tumorigenesis and provide a proof of mechanism for anti-cancer activity of VEGF inhibitors in this model.

Keywords: Lung cancer, Kinase and phosphatase inhibitors, Tumor angiogenesis, GEMM, VEGF

Background

Angiogenesis, formation of new blood vessels from existing vasculature, is an important process that supplies required nutrients and oxygen to cells which are distant from existing blood vessels. Angiogenesis is proven to play a key role in tumor growth and progression and several angiogenic factors such as VEGF (vascular endothelial growth factor), PDGF (platelet derived growth factor), bFGF (basic fibroblast growth factor) and HGF (hepatocyte growth factor) found to be among key regulators of tumor angiogenesis [1].

Series of investigations demonstrate a key role for VEGF in physiological or pathological angiogenesis [1]. Hence, a number of anti-angiogenic drugs targeting VEGF signaling pathway (ligand or the receptors) have been developed and are currently in use in cancer therapy. Bevacizumab (an anti-VEGF monoclonal antibody) was the first angiogenic inhibitor (hereafter AI) initially approved for use in patients with NSCLC (non-small cell lung cancer) or mCRC (metastatic colorectal cancer) [2,3]. Small molecule inhibitors of receptor tyrosine kinase inhibitors (RTKIs) are another class of agent targeting VEGF signaling pathway. RTKIs such as sunitinib, sorafenib, cediranib, motesanib, pazopanib and axitinib have been approved or are being

* Correspondence: farbodshojaei@hotmail.com

[†]Equal contributors

Oncology Research Unit, Pfizer Inc, La Jolla, CA 92121, USA

tested in different phases of clinical trials. Sunitinib which is a multi-targeted kinase inhibitor targets VEGFRs, C-SF1R, KIT and also platelet-derived growth factor (PDGFR) which plays an important role in blood vessel maturation [4]. Recently, sunitinib was approved by FDA for the treatment of advanced renal cell carcinoma, gastrointestinal stromal tumors and pancreatic neuroendocrine tumors [5,6]. Axitinib (AG-013736; Pfizer) is another oral potent tyrosine kinase inhibitor which mainly targets VEGFR and was approved by FDA for use in patients with advanced RCC [7]. In a murine lewis lung carcinoma model, single agent axitinib induced tumor necrosis and reduced microvessel density [8]. PF-00337210 (hereafter PF-210) is an oral, potent ATP-competitive inhibitor of VEGFR family [9]. It inhibits VEGFR2 phosphorylation and has greater selectivity towards VEGFR2 than other kinases. PF-210 has been shown to inhibit HUVEC cell survival *in vitro* and suppresses tumor angiogenesis in xenograft models [10].

Ras superfamily of proteins regulates cell growth, survival, and differentiation. Hras, Kras 4a, Kras 4b and Nras are the four highly homologous proteins encoded by three *Ras* genes [11,12]. Mutations in the *KRAS* gene lead to KRas protein activation in many human tumors including NSCLC, pancreatic cancer and colorectal cancer [12-14]. The majority of *KRAS* mutations (approximately 97%) occur in exon 2 at codon 12 and/or codon 13 in NSCLC patients [12]. The most common mutation in *KRAS* occurs at position 12, where glycine is replaced by a residue with side chain. NSCLC patients represent the majority of all lung cancer patients and remain a major cause of death [12]. Hence, Kras^{G12D-LSL} GEMM (genetically engineered mouse model) is one of the most relevant models of NSCLC to study tumor progression and to investigate efficacy of anti-cancer agents.

In the present study we investigated anti-tumor efficacy of three RTKIs including sunitinib, axitinib and PF-210 in Kras^{G12D-LSL} lung tumor model. Irrespective of the type (multi-targeted or selective), all three inhibitors significantly inhibited growth of advanced (adenocarcinoma) lesions in the lung indicating that VEGF is a key regulator of tumor angiogenesis in this model.

Methods

Tumor development and treatment in Kras^{G12D-LSL} GEMMs
Kras^{G12D-LSL} heterozygous mice were obtained from Jackson Laboratories (Jax West, CA) at approximately 3–4 weeks of age and were maintained by Pfizer La Jolla comparative medicine under guidelines provided by IACUC (Institutional Animal Care and Use Committee). Lung tumors were generated in Kras^{G12D-LSL} mice, using a recently published protocol [15]. Briefly, adenovirus expressing Cre recombinase (Adeno-Cre; the University

of Iowa Gene Transfer Vector Core, Iowa, IA) were titrated by Adenoviral Titration Kit (Clontech, CA) using instruction provided by the manufacturer. Prior to administration, Adeno-Cre virus was prepared in 50 μ l of plain MEM (minimal essential media; Gibco BRL; life Sciences, CA) supplemented with CaCl₂ (10 mM) followed by incubation at room temperature (RT) for 20 minutes. The recipients (n = 10) were anesthetized using Ketamine (Baxter) and Xylazine (Vedco) and the adeno-Cre preparation (2.5×10^7 infectious units; IU) was administered intra-nasally. To monitor tumor formation and progression, lung tissue was isolated (n = 1-3) at several time points (4, 8 and 12 wks) post inhalation and were stained with H&E (Hematoxylin and Eosin) using standard protocols in the laboratory [15]. The inhaled mice were randomized at 14 wks post-inhalation and were treated with vehicle, sunitinib (40 mg/kg qd), axitinib (15 mg/kg bid) and PF-210 (40 mg/kg qd) using oral route of administration and formulation protocols as described previously [8]. All the animal study procedures were monitored by the veterinary personnel to comply with guidelines provided by IACUC.

To assess therapeutic response to angiogenic inhibitors, lung lesions were quantified in the recipients by a certified pathologist. As previously described, lesions were categorized as hyperplastic, benign adenoma and adenocarcinoma [15]. Lesion quantification provided two types of analyses in the recipients: 1) percentage of each type of lesion in the recipient lung; 2) percentage of mice carrying these lesions in each treatment. To provide statistical analyses, we applied student's *t* test ($p < 0.05$ considered significant) to compare data between the vehicle *vs.* each treatment.

Histology

Formalin fixed paraffin embedded lung tissues were cut into 5 μ m sections and were stained for CD31, desmin, and F4-80 separately. Immunohistochemical staining was performed on Leica Bond III automated machine. Bond polymer refine detection kit was used for desmin and CD31 staining and bond intense R detection was used for F4-80 staining. For CD31 staining, lung sections were incubated for 45 minutes with rabbit anti-CD31 monoclonal antibody (clone SP38, Spring Bioscience, cat # M3384, 1:100 dilution). Desmin was stained by incubating lung section with mouse anti-huDesmin antibody (Dako Cytomation, cat# M0760, 1:1500 dilution) for 15 minutes. VEGFR1 and VEGFR2 was stained using anti-VEGFR1 antibody (abcam, cat# ab2350, 1:400 dilution) and anti-VEGFR2 antibody (cell signaling, cat# 2479, 1:200 dilution) respectively. Finally, F4-80 was stained with biotin anti-mouse F4-80 antibody (eBioscience, Cat # 13-4801-82, 1:75 dilution and


45 minutes incubation at RT). Images of stained-slides were captured using a Nanozoomer instrument (Hamamatsu, Japan) and the data was analyzed using Aperio Imagescope software.

Results

Targeting the VEGF pathway is sufficient to inhibit progression of lung adenocarcinoma lesions in $Kras^{G12D-LSL}$ mice

Our strategy to investigate anti-tumor efficacy of AIs in $Kras^{G12D-LSL}$ mice is depicted in Figure 1A. $Kras^{G12D-LSL}$ mice were inhaled intranasally with Adeno-Cre at 6–8 weeks of age and were maintained without any further intervention. At 8–10 weeks post inhalation, few mice ($n = 1-3$) were randomly euthanized to assess tumor

formation and progression in the lung (Figure 1B). All the remaining mice were randomized into treatment groups and treated with vehicle, PF-210, axitinib and sunitinib for about 8 weeks. Upon termination of the study, lung tissues were analyzed for tumor lesions using H&E staining (Figure 2). Compared to vehicle-treated group, there was a significant reduction in lung lesion in all the three drugs treated groups. To further understand mechanism of action of AIs, we classified lung lesions into three categories including hyperplasia, benign neoplasia and malignant (adenocarcinoma). Detailed pathology analyses of lesions revealed that hyperplastic lesions were not significantly affected by AIs compared to control-treated animals (Figure 3A). However, the percentage of benign neoplastic lesions was significantly ($p < 0.01$) inhibited by PF-210 (more







Figure 2 AIs suppress growth of advanced lesions in lung tumors in *Kras*^{G12D-LSL} mice. As described in Figure 1, *Kras*^{G12D-LSL} tumor bearing mice were treated for 8 wks with AIs. Treatments started at 14 wks post administration of Adeno-Cre. At the end of study, all the lung tissues were isolated from the recipients and were analyzed using H&E staining. Images are the representative of each treatment.

than 90%) and also axitinib or sunitinib (more than 50%) compared to vehicle-treated mice. Finally malignant lesions were significantly ($p < 0.01$) inhibited by all the AIs. Additionally we investigated percentage of mice carrying the above-mentioned lesions (Figure 3B). Irrespective of the type of treatment, all mice carried hyperplastic lesions. While all mice treated with axitinib or sunitinib carried benign neoplasia, only 40% of PF-210 treated animals carried these lesions indicating the potency of this compound. Finally all three AIs reduced frequency of malignant lesions by at least 50% in treated mice (Figure 3B). Overall, two types of analyses (percentage of lesions in the lung and percentage of mice carrying specific lesion) indicate that AIs specifically target advanced lesions (malignant or adenocarcinoma).

Components of vasculature and stroma are targeted by AIs

To further investigate tumor vasculature, we stained lung tissues with different markers such as CD31 and desmin to stain endothelial cells and smooth muscle

cells respectively [16]. Vasculature analysis by CD31 staining showed high density of tumor blood vessels in adenoma and adenocarcinoma lesions in the vehicle group (Figure 4). Moreover, these vessels were desmin positive indicative of a mature vasculature in these lesions. In contrast, tumor lesions in AI treated groups had less number of blood vessels further suggesting that vasculature is the main target of these AIs. Additionally, vasculature was found to be more fragmented compared to the blood vessels in vehicle treated mice. Similar to CD31 staining, all three AIs (especially PF-210 and axitinib) targeted smooth muscle cells suggesting that not only blood vessels but also other components of vasculature are affected. We also investigated the effects of AIs on the expression of VEGFR1 and VEGFR2 which play an important role in angiogenesis and tumor progression [1]. High levels of VEGFR1 was observed on tumor cells in vehicle treated mice (Figure 4) that is consistent with the expression of VEGFR1 on tumor cells isolated from *Kras* mutant NSCLC tumors in an earlier report [17].



Interestingly, sunitinib and PF-210, but not axitinib, inhibited VEGFR1 expression on tumor cells (Figure 4). Compared to vehicle treated tumors that expressed abundant levels of VEGFR2 on blood vessels, all three AIs inhibited VEGFR2 expression on the tumor vasculature further providing a mechanism for the anti-angiogenic activity of these compounds. Overall, these results suggest that inhibition of angiogenesis is the main mechanism by which AIs suppress growth of benign and malignant lesions in this model of NSCLC.

Tumor associated macrophages (TAMs) are a key component of tumor microenvironment and have been implicated in tumor progression and angiogenesis. It has been shown that NSCLC patients with higher density of TAMs have lower median relapse-free survival (7 months) compared to patients whose tumors had lower density of TAMs (26 months survival) [18]. Macrophage staining (using F4-80) indicated infiltration of these TAMs in the lung in vehicle treated mice (Figure 4). Treatment with AIs particularly sunitinib and axitinib was associated with lower density of TAMs further suggesting an additional mechanism for anti-tumor efficacy of AIs in $Kras^{G12D-LSL}$ lung tumors.

Discussion

This study reports anti-tumor efficacy of three different RTKIs including PF-210, axitinib and sunitinib in spontaneous tumors in lung in $Kras^{G12D-LSL}$ GEMMs. The high failure rate of clinical trials in late stage cancer patients warrants development of mouse tumor models which are more relevant to the human diseases. GEMMs, carrying genetic alterations similar to what is observed in cancer patients, might represent a more relevant tumor model to predict clinical outcome.

The VEGF signaling pathway is one of the major signaling pathways in tumor angiogenesis in many cancers. An anti-VEGF monoclonal antibody, bevacizumab, has been approved in combination with chemotherapy for the treatment of NSCLC [19]. Bevacizumab is the first targeted agent to improve survival in advanced-stage NSCLC patients when combined with first-line chemotherapy. In the present study, we use sunitinib, axitinib, PF-210 all of which targeting VEGFR signaling pathway with different pharmacokinetic and pharmacodynamic properties [7,20]. Our results show that reduction of malignant lesions in lungs is the common and consistent theme among all the above compounds. Progression of malignant lesions prior to diagnosis and treatment are the major contributors to low survival rate in NSCLC patients [18]. Lack of efficacy of these agents in hyperplastic lesions indicate that angiogenesis may not play a significant role in growth of pre-neoplastic lesions lung tumors in $Kras^{G12D-LSL}$ GEMMs. Additionally while sunitinib is a multi-targeting RTKIs, our data indicate that, at clinical dose, targeting PDGFR- β , KIT and CSF1-R does not provide additional efficacy compared to PF-210 and axitinib which are selective inhibitors of VEGF. These data once again signifies the role of VEGF as a key regulator of tumor angiogenesis in a preclinical model of NSCLC. PF-210 showed superior efficacy in suppressing benign neoplasia lesions compared to axitinib and sunitinib. Future investigations




Figure 4 Tumor vasculature and components of stroma in lung tumors are targeted by anti-angiogenesis. Formalin fixed paraffin embedded lung sections (5 μ m) stained with anti-CD31 (endothelial cells), anti-desmin (smooth muscle cells), anti-F4-80 (macrophages), anti-VEGFR1 and anti-VEGFR2 as described in the materials and methods. The slides were scanned on Nanozoomer instrument using 20 \times magnification. Irrespective of treatment, AIs treated lungs showed lower CD31 and desmin density. F4-80 staining in vehicle treated mice showed greater infiltration of macrophages. However, AIs particularly axitinib and sunitinib significantly reduced macrophage infiltration in the tumors. AIs had differential effects on VEGFR1 and VEGFR2 expression since all treatments reduced VEGFR2 expression in the tumor vasculature, VEGFR1 expression was mainly inhibited by PF-210 and sunitinib. Pictures are the representative images of each staining from each treatment.

might provide some insight into the mechanism of action of PF-210.

Histopathological analysis showed that all these AIs target tumor vasculature to inhibit growth of malignant lesions. Moreover, most of the tumor blood vessels in treated mice lacked smooth muscle cell coverage suggesting a role for VEGF in establishment of a cross talk between smooth muscle cells and endothelial cells. Furthermore, AI- treated mice had lower number of TAMs compared to the vehicle treated animals suggesting that these cells may play a proangiogenic role in this model [21]. Future studies will determine if AIs alter homing of macrophages to the tumors or are directly targeting them. In addition, further investigation is warranted to understand pharmacokinetics and

pharmacodynamics of these compounds in the tumors which may describe differences in the mechanism of action of AIs in the current study.

Conclusion

Our data indicate that small molecule inhibitors of VEGF pathway suppress growth of adenocarcinoma lesions in a NSCLC model of $Kras^{G12D-LSL}$ GEMM by targeting components of tumor vasculature and stroma.

Competing interests

All authors are employees and are stock holders of Pfizer Incorportations.

Authors' contributions

FS designed all the studies and experiments, BKM performed all the histology staining and analyzed the data, JHL, BHS and FS executed in vivo study, BKM and FS wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We would like to thank Pamela Vizcarra for assisting in histology staining and James G Christensen for helpful discussion during preparation of the manuscript.

Received: 3 October 2012 Accepted: 25 April 2013

Published: 29 April 2013

References

1. Shojaei F: **Anti-angiogenesis therapy in cancer: current challenges and future perspectives.** *Cancer Lett* 2012, **320**(2):130–137.
2. Sandler A, Gray R, Perry MC, Brahmer J, Schiller JH, Dowlati A, Lilenbaum R, Johnson DH: **Paclitaxel-carboplatin alone or with bevacizumab for non-small-cell lung cancer.** *N Engl J Med* 2006, **355**(24):2542–2550.
3. Zondor SD, Medina PJ: **Bevacizumab: an angiogenesis inhibitor with efficacy in colorectal and other malignancies.** *Ann Pharmacother* 2004, **38**(7–8):1258–1264.
4. Sun L, Liang C, Shirazian S, Zhou Y, Miller T, Cui J, Fukuda JY, Chu JY, Nematalla A, Wang X, et al: **Discovery of 5-[5-fluoro-2-oxo-1,2-dihydroindol-(3Z)-ylidenemethyl]-2,4-dimethyl-1H-pyrrole-3-carboxylic acid (2-diethylaminoethyl) amide, a novel tyrosine kinase inhibitor targeting vascular endothelial and platelet-derived growth factor receptor tyrosine kinase.** *J Med Chem* 2003, **46**(7):1116–1119.
5. Faivre S, Demetri G, Sargent W, Raymond E: **Molecular basis for sunitinib efficacy and future clinical development.** *Nat Rev Drug Discov* 2007, **6**(9):734–745.
6. Goodman VL, Rock EP, Dagher R, Ramchandani RP, Abraham S, Gobburu JV, Booth BP, Verbois SL, Morse DE, Liang CY, et al: **Approval summary: sunitinib for the treatment of imatinib refractory or intolerant gastrointestinal stromal tumors and advanced renal cell carcinoma.** *Clin Cancer Res* 2007, **13**(5):1367–1373.
7. Rugo HS, Herbst RS, Liu G, Park JW, Kies MS, Steinfeldt HM, Pithavala YK, Reich SD, Freddo JL, Wilding G: **Phase I trial of the oral antiangiogenesis agent AG-013736 in patients with advanced solid tumors: pharmacokinetic and clinical results.** *J Clin Oncol* 2005, **23**(24):5474–5483.
8. Hu-Lowe DD, Zou HY, Grazzini ML, Hallin ME, Wickman GR, Amundson K, Chen JH, Rewolinski DA, Yamazaki S, Wu EY, et al: **Nonclinical antiangiogenesis and antitumor activities of axitinib (AG-013736), an oral, potent, and selective inhibitor of vascular endothelial growth factor receptor tyrosine kinases 1, 2, 3.** *Clin Cancer Res* 2008, **14**(22):7272–7283.
9. Marrone TJ H-LD, Grazzini M, Yin MJ, Chen J, Hallin M, et al: **PF-00337210, a potent, selective and orally bioavailable small molecule inhibitor of VEGFR-2.** In *Proceedings of the 98th Annual Meeting of the American Association for Cancer Research; AACR Meeting Abstracts: 2007.* Los Angeles, CA: AACR; 2007:3992.
10. Liu: **Phase I pharmacokinetic (PK) and pharmacodynamic (PD) study of PF-00337210, a highly selective VEGFR inhibitor.** *J Clin Oncol* 2009, **27**:3519.
11. Chang EH, Gonda MA, Ellis RW, Scolnick EM, Lowy DR: **Human genome contains four genes homologous to transforming genes of Harvey and Kirsten murine sarcoma viruses.** *Proc Natl Acad Sci U S A* 1982, **79**(16):4848–4852.
12. Riely GJ, Marks J, Pao W: **KRAS mutations in non-small cell lung cancer.** *Proc Am Thorac Soc* 2009, **6**(2):201–205.
13. Hingorani SR, Petricoin EF, Maitra A, Rajapakse V, King C, Jacobetz MA, Ross S, Conrads TP, Veenstra TD, Hitt BA, et al: **Preinvasive and invasive ductal pancreatic cancer and its early detection in the mouse.** *Cancer Cell* 2003, **4**(6):437–450.
14. Aguirre AJ, Bardeesy N, Sinha M, Lopez L, Tuveson DA, Horner J, Redston MS, DePinho RA: **Activated Kras and Ink4a/Arf deficiency cooperate to produce metastatic pancreatic ductal adenocarcinoma.** *Genes Dev* 2003, **17**(24):3112–3126.
15. Simmons BH, Lee JH, Lalwani K, Giddabasappa A, Snider BA, Wong A, Lappin PB, Eswaraka J, Kan JL, Christensen JG, et al: **Combination of a MEK inhibitor at sub-MTD with a PI3k/mTOR inhibitor significantly suppresses**

growth of lung adenocarcinoma tumors in Kras(G12D-LSL) mice. *Cancer Chemother Pharmacol* 2012, **70**(2):213–220.

16. Pollock L, Rampling D, Greenwald SE, Malone M: **Desmin expression in rhabdomyosarcoma: influence of the desmin clone and immunohistochemical method.** *J Clin Pathol* 1995, **48**(6):535–538.
17. Roybal JD, Zang Y, Ahn YH, Yang Y, Gibbons DL, Baird BN, Alvarez C, Thilaganathan N, Liu DD, Saintigny P, et al: **miR-200 Inhibits lung adenocarcinoma cell invasion and metastasis by targeting Flt1/VEGFR1.** *Mol Canc Res* 2011, **9**(1):25–35.
18. Chen JJ, Lin YC, Yao PL, Yuan A, Chen HY, Shun CT, Tsai MF, Chen CH, Yang PC: **Tumor-associated macrophages: the double-edged sword in cancer progression.** *J Clin Oncol* 2005, **23**(5):953–964.
19. Planchard D: **Bevacizumab in non-small-cell lung cancer: a review.** *Expert Rev Anticancer Ther* 2011, **11**(8):1163–1179.
20. Faivre S, Delbaldo C, Vera K, Robert C, Lozahic S, Lassau N, Bello C, Deprimo S, Brega N, Massimini G, et al: **Safety, pharmacokinetic, and antitumor activity of SU11248, a novel oral multitarget tyrosine kinase inhibitor, in patients with cancer.** *J Clin Oncol* 2006, **24**(1):25–35.
21. Allavena P, Mantovani A: **Immunology in the clinic review series; focus on cancer: tumour-associated macrophages: undisputed stars of the inflammatory tumour microenvironment.** *Clin Exp Immunol* 2012, **167**(2):195–205.

doi:10.1186/1471-2407-13-213

Cite this article as: Majeti et al.: VEGF is an important mediator of tumor angiogenesis in malignant lesions in a genetically engineered mouse model of lung adenocarcinoma. *BMC Cancer* 2013 **13**:213.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

