

Research article

Open Access

Both *SEPT2* and *MLL* are down-regulated in *MLL-SEPT2* therapy-related myeloid neoplasia

Nuno Cerveira¹, Joana Santos¹, Susana Bizarro¹, Vera Costa¹,
Franclim R Ribeiro¹, Susana Lisboa¹, Cecília Correia¹, Lurdes Torres¹,
Joana Vieira¹, Simone Snijder², José M Mariz³, Lucília Norton⁴,
Clemens H Mellink², Arjan Buijs⁵ and Manuel R Teixeira^{* 1,6}

Address: ¹Department of Genetics, Portuguese Oncology Institute, Porto, Portugal, ²Department of Clinical Genetics, Academic Medical Center, Amsterdam, The Netherlands, ³Department of Hemato-Oncology, Portuguese Oncology Institute, Porto, Portugal, ⁴Department of Pediatrics, Portuguese Oncology Institute, Porto, Portugal, ⁵Department of Medical Genetics, University Medical Center, Utrecht, The Netherlands and ⁶Abel Salazar Biomedical Sciences Institute (ICBAS), Porto, Portugal

Email: Nuno Cerveira - nscerveira@gmail.com; Joana Santos - joana.cris.santos@gmail.com; Susana Bizarro - susanabizarro@gmail.com; Vera Costa - veralmcosta@gmail.com; Franclim R Ribeiro - ribeirofr@gmail.com; Susana Lisboa - susanalisboa@hotmail.com; Cecília Correia - correia.cecilia@gmail.com; Lurdes Torres - lurdes_torres@hotmail.com; Joana Vieira - joana_vieira@netcabo.pt; Simone Snijder - s.snijder@amc.uva.nl; José M Mariz - mariz@ipopoporto.min-saude.pt; Lucília Norton - lucilia.norton@ipopoporto.min-saude.pt; Clemens H Mellink - c.h.mellink@amc.uva.nl; Arjan Buijs - a.buijs@umcutrecht.nl; Manuel R Teixeira* - manuel.teixeira@ipopoporto.min-saude.pt

* Corresponding author

Published: 15 May 2009

Received: 3 December 2008

BMC Cancer 2009, 9:147 doi:10.1186/1471-2407-9-147

Accepted: 15 May 2009

This article is available from: <http://www.biomedcentral.com/1471-2407/9/147>

© 2009 Cerveira et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: A relevant role of septins in leukemogenesis has been uncovered by their involvement as fusion partners in *MLL*-related leukemia. Recently, we have established the *MLL-SEPT2* gene fusion as the molecular abnormality subjacent to the translocation t(2;11)(q37;q23) in therapy-related acute myeloid leukemia. In this work we quantified *MLL* and *SEPT2* gene expression in 58 acute myeloid leukemia patients selected to represent the major AML genetic subgroups, as well as in all three cases of *MLL-SEPT2*-associated myeloid neoplasms so far described in the literature.

Methods: Cytogenetics, fluorescence in situ hybridization (FISH) and molecular studies (RT-PCR, qRT-PCR and qMSP) were used to characterize 58 acute myeloid leukemia patients (AML) at diagnosis selected to represent the major AML genetic subgroups: *CBFB-MYH11* (n = 13), *PML-RARA* (n = 12); *RUNX1-RUNX1T1* (n = 12), normal karyotype (n = 11), and *MLL* gene fusions other than *MLL-SEPT2* (n = 10). We also studied all three *MLL-SEPT2* myeloid neoplasia cases reported in the literature, namely two AML patients and a t-MDS patient.

Results: When compared with normal controls, we found a 12.8-fold reduction of wild-type *SEPT2* and *MLL-SEPT2* combined expression in cases with the *MLL-SEPT2* gene fusion (p = 0.007), which is accompanied by a 12.4-fold down-regulation of wild-type *MLL* and *MLL-SEPT2* combined expression (p = 0.028). The down-regulation of *SEPT2* in *MLL-SEPT2* myeloid neoplasias was statistically significant when compared with all other leukemia genetic subgroups (including those with other *MLL* gene fusions). In addition, *MLL* expression was also down-regulated in the group of *MLL* fusions other than *MLL-SEPT2*, when compared with the normal control group (p = 0.023)

Conclusion: We found a significant down-regulation of both *SEPT2* and *MLL* in *MLL-SEPT2* myeloid neoplasias. In addition, we also found that *MLL* is under-expressed in AML patients with *MLL* fusions other than *MLL-SEPT2*.

Background

Septins comprise an evolutionarily conserved family of GTP-binding proteins that are found primarily in fungi and animals [1]. In humans, 14 septin genes have been characterized to date (*SEPT1* to *SEPT14*). All septin transcripts contain multiple translation initiation sites and are alternatively spliced, giving origin to multiple septin isoforms, some of which are tissue specific [1]. Although the precise functions of septins remain unclear, current data suggest that they coordinate changes in cytoskeletal and membrane organization by acting as scaffolds that recruit factors to precise sites in a cell and/or as barriers that segregate membrane areas into discrete domains [1,2]. For instance, the human *SEPT2* associates with *SEPT6* and *SEPT7* to form an hexamer complex that is the core unit for generation of septin filaments associated with the contractile ring in dividing cells, being therefore essential for proper cytokinesis and chromosome segregation [2,3].

Septins have been reported to be deregulated in various human diseases, including cancer [4]. A relevant role of septins in leukemogenesis has been uncovered by their involvement as fusion partners in *MLL*-related leukemia. We have established the *MLL-SEPT2* gene fusion as the molecular abnormality subjacent to the translocation t(2;11)(q37;q23) in therapy-related acute myeloid leukemia (t-AML) [5]. Subsequently, van Binsbergen et al [6] identified a second *MLL-SEPT2* fusion variant in a patient with t-AML and we have recently uncovered a third *MLL-SEPT2* alternative fusion variant in a case of therapy-related myelodysplastic syndrome (t-MDS) [7,8]. Four other septin family genes (*SEPT5*, *SEPT6*, *SEPT9* and *SEPT11*) had previously been identified as *MLL* fusion partners in leukemia, making the septins the protein family with more numbers involved in *MLL*-related leukemia [9-13], and suggesting that their involvement in leukemogenesis is not a chance event. This hypothesis is supported by the fact that all the reported *MLL*-septin fusions are in frame and the breakpoints are found at the very 5' end of known septin open reading frames [5,9-13]. In this work we show evidence that the fusion of *MLL* with *SEPT2* is associated with down-regulation of both *SEPT2* and *MLL* expression in t-AML/t-MDS.

Methods

Patient samples

We studied 58 acute myeloid leukemia patients (AML) at diagnosis selected to represent the major AML genetic subgroups, including 13 cases with a *CBFB-MYH11* rearrangement, 12 cases with a *PML-RARA* rearrangement, 12 cases with a *RUNX1-RUNX1T1* rearrangement, 11 cases with normal karyotype, and 10 cases with rearrangements of the *MLL* gene other than *MLL-SEPT2* [see Additional file 1]. We also studied all three *MLL-SEPT2* patients reported in the literature, namely two with t-AML (patients 59 and 60) and the third with t-MDS (patient 61), which were the

primary targets of our investigation [see Additional file 1] [5-8]. All but patients 60 and 61 were treated at the Portuguese Oncology Institute, Porto, Portugal, and bone marrow samples were obtained to perform cytogenetic and molecular studies. Patients 60 and 61 were treated in the Netherlands at the St. Antonius Hospital, Nieuwegein [6] and the Amsterdam Academic Medical Center [7], respectively, from whom cDNA was obtained.

As a control group we studied bone marrow samples obtained from ten individuals studied for the purpose to rule out a hematological disease.

This study was approved by the Portuguese Oncology Institute ethic committee, and informed consent was obtained from all patients.

Chromosome banding and molecular cytogenetics

The diagnostic bone marrow samples of the Portuguese patients (cases 1 to 59) were cultured for 24 hours in RPMI 1640 medium with GlutaMAX-I (Invitrogen, London, UK) supplemented with 20% fetal bovine serum (Invitrogen, London, UK). Chromosome preparations were made by standard methods and banded by trypsin-Leishman. Karyotypes were described according to the International System for Human Cytogenetic Nomenclature [14].

Whenever appropriate, fluorescence in situ hybridization (FISH) analysis for specific fusion genes or rearrangements was performed using dual-color, break-apart or dual-fusion, probes (Vysis, Downers Grove, USA).

Chromosome banding and molecular cytogenetic analyses of patients 60 and 61 were described previously [6,7].

RNA extraction and cDNA synthesis

Total RNA was extracted from the diagnostic bone marrow sample of patients 1 to 59 and controls using 1 ml of Tripure isolation reagent (Roche Diagnostics, Indianapolis, USA) and quantified in a NanoDrop ND-100 spectrophotometer (NanoDrop Technologies, Wilmington, USA). For cDNA synthesis, 1 µg of total RNA was subjected to reverse transcription with random hexamers using the Superscript III First-Strand Synthesis System for RT-PCR (Invitrogen, Carlsbad, USA), according to the manufacturer's instructions. The final cDNA was diluted with 30 µl of H₂O. cDNA quantity and quality were assessed in a NanoDrop ND-100 spectrophotometer (NanoDrop Technologies, Wilmington, USA).

Qualitative Reverse-Transcription Polymerase Chain Reaction (RT-PCR)

RT-PCR assays for detection of the fusion transcripts *RUNX1-RUNX1T1*, *CBFB-MYH11*, and *PML-RARA* were performed on the diagnostic samples according to the

BIOMED-1 protocol [15]. The primers and PCR reaction conditions for the detection of rearrangements involving the *MLL* gene were previously published [5,10,16-19].

Quantitative Real-Time Polymerase Chain Reaction (qRT-PCR)

We have evaluated the mRNA expression of *MLL* and *SEPT2* genes by qRT-PCR on an ABI PRISM 7000 Sequence Detection System (Applied Biosystems, Foster City, USA). Primers and probes for *MLL* and *SEPT2* were derived from the published mRNA sequences of *SEPT2* and *MLL* (GenBank accession nos. [NM_001008491.1](#) and [NM_005933.2](#), respectively), and designed with Primer Express 2.0 (Applied BioSystems) and purchased from Metabion (Metabion, Martinsried, Deutschland) [see Additional file 2]. Primers and probes for the *ABL1* gene (GenBank accession no. [NM_005157](#)), used as endogenous control, were previously described and approved for qRT-PCR based diagnosis and minimal residual disease detection in leukaemic patients, due to be similarly expressed in normal and diagnostic samples as well as within normal samples [see Additional file 2] [20,21]. All primers and probes were designed outside of *MLL* and *SEPT2* breakpoint cluster regions in exons 4–5 and exons 3–4, respectively. To determine the relative expression levels of the target gene in each sample, the relative amount of the target gene was calibrated to the relative amount of the internal reference gene and expressed in terms of ratios between the target and the reference that were then multiplied by 100 for easier tabulation (target gene/*ABL1* × 100). PCR reactions were performed in a 25 µl volume containing 5 µl of synthesized cDNA, 12.5 µl of TaqMan universal PCR master mix, 0.3 µM of each primer and 0.2 µM of each probe. PCR was performed in separate wells for each primer/probe set and each sample was run in triplicate. PCR parameters were as follows: 50°C for 2 min., 95°C for 10 min., followed by 50 cycles at 95°C for 15 s. and 60°C for 1 min. Each plate included non-template controls and serial dilutions of a strongly expressing sample (*MLL* or *SEPT2*) to construct the standard curves.

Bisulfite treatment

Sodium bisulfite conversion of unmethylated (but not methylated) cytosine residues to uracil in a sample of genomic DNA obtained from *MLL-SEPT2* case 59 (DNA from *MLL-SEPT2* cases 60 and 61 was not available) and three normal controls was performed as previously described [22]. Briefly, 500 ng of genomic DNA was denatured with 0.3 M NaOH in a total volume of 21 µl for 20 min. at 50°C. A volume of 450 µl freshly prepared bisulfite solution (2.5 M sodium bisulfite, 125 mM hydroquinone, and 0.2 M NaOH) was added to each denaturation reaction, and the mixture was incubated at 70°C for 3 hours in the dark. The resulting bisulfite-converted DNA was then purified by using Wizard DNA purification resin

(Wizard DNA Clean-Up System; Promega, Madison, USA) according to the manufacturer's instructions and eluted in 45 µl of water preheated at 70°C. The eluted DNA was denatured in 0.3 M NaOH for 10 min. at room temperature. Finally, the bisulfite converted and denatured genomic DNA was precipitated with 100% ethanol, dried, resuspended in 30 µl of water, and stored at -20°C.

Quantitative Methylation-Specific Polymerase Chain Reaction (qMSP)

Due to the very low quantity of DNA available from *MLL-SEPT2* case 59, only the *SEPT2* gene could be analyzed by qMSP. The *SEPT2* 5'-CpG island was identified using the CpG Island Searcher <http://cpgislands.com> [23] and the *SEPT2* genomic sequence (GenBank accession no. [NT_005416.12](#)). That CpG island was found to encompass the predicted *SEPT2* promoter region using PROSCAN 1.7 [24]. Primers and probe for the *SEPT2* 5'-CpG island were designed with Methyl Primer Express 1.0 (Applied BioSystems, Foster City, USA), and selected to specifically amplify fully methylated bisulfite-converted DNA [see Additional file 2]. Primers and probe for the internal reference gene, *ACTB* (GenBank accession no. [NM_001101](#)), were described previously [25] and were designed to amplify and detect a region of the gene that is devoid of CpG nucleotides to normalize for DNA input in each sample [see Additional file 2]. qMSP of the chemically modified DNA was performed in an ABI PRISM 7000 Sequence Detection System (Applied Biosystems, Foster City, USA), as previously described [26]. In brief, fluorescence based real-time PCR assays were carried out in a reaction volume of 20 µl, consisting of 16.6 mM ammonium sulfate, 67 mM trizma preset, 6.7 mM MgCl₂, 10 mM mercaptoethanol, 0.1% DMSO, 200 µM each of dATP, dCTP, dGTP, and dTTP, 600 nM of each primer, 0.4 µl of Rox dye, 200 nM of probe, 1 unit of platinum Taq polymerase (Invitrogen, Carlsbad, USA), and 2 µl of bisulfite-modified DNA as a template. PCR was performed in separate wells for each primer/probe set and each sample was run in triplicate. PCR was performed under the following conditions: 95°C for 2 min., followed by 45 cycles of 95°C for 15 s. and 60°C for 1 min.

To ensure the specificity of the analysis, each 96-well PCR plate had wells that contained completely methylated DNA at all CpGs (positive control – CpGenome Universal Methylated DNA, Chemicon Europe, Hampshire, UK), a completely unmethylated DNA (negative control – CpGenome Universal Unmethylated DNA, Chemicon Europe, Hampshire, UK), and multiple water blanks (contamination control). To determine the relative levels of methylated promoter DNA in each sample, we used serial dilutions of the positive control DNA to construct the calibration curve. The values obtained (mean quantity) for each target gene were divided by the respective values of

the internal reference gene (*ACTB*). The ratio thus generated, which constitutes an index of the percentage of input copies of DNA that are fully methylated at the primer- and probe-binding sites, was then multiplied by 100 for easier tabulation (methylation level = target gene/reference gene × 100).

Statistical analyses

Normalized expression values for *MLL* and *SEPT2* were compared among the different sample groups using the non-parametric Kruskal-Wallis H and Mann-Whitney U tests. The correlation between *MLL* and *SEPT2* expression values within each group was assessed using Pearson's test. All analyses were performed using SPSS version 15.0 (SPSS, Chicago, USA).

Results

In all cases, RNA and/or cDNA quantity and quality was evaluated and was found to be appropriate for expression studies. The 260/280 and 260/230 absorbance ratios for RNA samples were in the range of 1.8–2.2. For the cDNA samples, the 260/280 and 260/230 sample absorbance ratios were in the range of 1.6–2.0 and 1.8–2.2, respectively.

Normalized expression levels for *MLL* and *SEPT2* within each sample group are depicted in Figure 1. Statistically significant differences were observed for the combined wild-type *SEPT2* and *MLL-SEPT2* expression in the *MLL-SEPT2* cases when compared with the normal controls, showing a 12.8-fold lower median expression in the *MLL-SEPT2* subset (p = 0.007). Furthermore, the combined wild-type *SEPT2* and *MLL-SEPT2* expression was significantly lower (5.4 to 9.4 fold) in the *MLL-SEPT2* cases than in all other leukemia genetic subgroups (Table 1). The combined expression of wild-type *MLL* and *MLL-SEPT2* was also significantly lower (12.4 fold; p = 0.028) in the *MLL-SEPT2* myeloid neoplasias when compared with the normal controls, as well as with the *CBFB-MYH11* and *RUNX1-RUNX1T1* leukemia subgroups (13.4 and 10.5 fold, respectively). We next investigated whether DNA hypermethylation of the 5' *SEPT2* region was contributing to the *SEPT2* gene down-regulation, by examining the methylation status of the CpG island located upstream of the *SEPT2* gene transcriptional initiation site [2225 base pair (bp) in length (-7457 to -9257)]. *SEPT2* 5' CpG island hypermethylation was detected in the positive control, but not in the *MLL-SEPT2* case 59 or the normal controls.

No statistically significant differences were observed for the wild-type *SEPT2* expression between the non-*MLL-SEPT2* leukemia subgroups and the normal controls, with the exception of the *PML-RARA* and "normal karyotype" leukemias that showed lower expression (Table 1). No statistically significant differences were observed for the




Figure 1
Box-plots of normalized expression values for *SEPT2* (A) and *MLL* (B) in subgroups of myeloid malignancies and control samples. Asterisks denote significant differences when compared with the normal control group. The combined expression of *MLL-SEPT2* and *MLL* or *SEPT2* was significantly lower in *MLL-SEPT2* patients compared to all other groups (see Table 1 for significance values).

wild-type *MLL* expression between the non-*MLL-SEPT2* leukemia subgroups and the normal controls, with the exception of the significantly lower expression seen in the patient group with *MLL* fusions with other partners other than *SEPT2* (q = 0,023).

Discussion

Fusion oncogenes are generally thought to contribute to carcinogenesis by either causing over-expression of the 3' partner due to promoter swap or by originating a chimeric

Table 1: Normalized values for the expression of MLL-SEPT2 and MLL or SEPT2 in subgroups of hematologic malignancies and normal controls

Genetic groups	n	SEPT2 + MLL-SEPT2				MLL + MLL-SEPT2			
		Median (P25-P75)	Fold Change (a)	P-value (a)	P-value (b)	Median (P25-P75)	Fold Change (a)	P-value (a)	P-value (b)
<i>CBFB-MYH11</i>	13	3388 (2411-5634)	9.4	0.004	Ns	2692 (1698-5099)	13.4	0.025	ns
<i>PML-RARA</i>	12	1957 (1055-2904)	5.4	0.004	0.001	1083 (971-2698)	5.4	ns	ns
<i>RUNX1-RUNX1T1</i>	12	3187 (2100-3760)	8.9	0.004	ns	2107 (1765-3018)	10.5	0.018	ns
Normal Karyotype	11	2242 (1604-3424)	6.2	0.005	0.010	1621 (1012-2895)	8.1	ns	ns
<i>MLL</i> Fusions	10	3069 (2159-4600)	8.5	0.007	ns	1292 (708-2050)	6.4	ns	0.023
<i>MLL-SEPT2</i>	3	359 (333-724)	1.0	-	0.007	200 (180-1612)	1.0	-	0.028
Normal Control	10	4599 (2939-6365)	12.8	0.007	-	2494 (1749-3929)	12.4	0.028	-

(a) Comparison of each group with *MLL-SEPT2* patients; (b) Comparison of each group with normal control group; (P25) percentile 25; (P75) percentile 75.

protein with new biochemical properties. Surprisingly, when compared with the normal controls, we found a 12.8-fold reduction of the combined *MLL-SEPT2* and wild-type *SEPT2* expression in myeloid neoplasias with the *MLL-SEPT2* gene fusion, which is accompanied by 12.4-fold down-regulation of the combined *MLL-SEPT2* and wild-type *MLL* expression. The down-regulation of *SEPT2* in *MLL-SEPT2* myeloid neoplasias was also statistically significant when compared with all other leukemia genetic subgroups (including those with other *MLL* gene fusions). It is conceivable that deregulation of *SEPT2* expression can occur as a result of its fusion with *MLL* (for example by haplo-insufficiency), since the *MLL-SEPT2* gene is under the control of the *MLL* promoter. However, not only the magnitude of *SEPT2* under-expression far exceeds the maximum 50% reduction that would be expected if one of the *SEPT2* gene copies has its expression shutdown, but wild-type *MLL* expression seems to be strongly down-regulated as well. This suggests a concomitant down-regulation of wild-type *MLL*, wild-type *SEPT2*, and the *MLL-SEPT2* fusion gene. The fact that *MLL*, *SEPT2*, and *MLL-SEPT2* map to distinct chromosomes excludes a localized transcriptional repression affecting contiguous genes via a long-range control element as a possible mechanism.

Interestingly, *MLL* expression was also down-regulated in the group of *MLL* fusions other than *MLL-SEPT2*, when compared with the normal control group. Wild-type *MLL* down-regulation associated with *MLL* abnormalities was previously observed in AML with *MLL* partial tandem duplication (PTD) [27]. In that instance, the wild-type

MLL transcript derived from the non-rearranged *MLL* allele was absent in the majority of cases of *MLL*-PTD, with the authors suggesting that the silencing of wild-type *MLL* may result from the action of the *MLL*-PTD protein via an auto-regulatory mechanism [27], which has so far not been described for *MLL-SEPT2*. In addition, down-regulation of *MLL* when fused with a partner gene was also previously observed in *MLL-MLLT3* patients [27], suggesting that this can be a common event in *MLL*-related leukemia. Since *MLL* fusion proteins seem to transform by a gain-of-function mechanism with conversion of the *MLL* chimera into a potent transcriptional activator [28,29], quantitative oscillations in wild-type and chimeric *MLL* expression level presumably do not abrogate the leukemogenic properties of *MLL* fusion proteins. One alternative explanation for the observed down-regulation of *MLL*, *SEPT2* and *MLL-SEPT2* is the involvement of a transcriptional rather than post-transcriptional mechanism, for instance regulated via an epigenetic mark. DNA methylation within the promoter region of a gene can result in chromatin compaction and inhibition or down-regulation of gene transcription, and aberrant promoter methylation in cis is often responsible for gene silencing in a variety of malignancies [30]. However, the absence of DNA methylation in the CpG island located 5' of the *SEPT2* gene (encompassing the predicted *SEPT2* promoter region) suggests that hypermethylation of wild-type *SEPT2* is probably not the mechanism responsible for the observed gene silencing but, since in only one *MLL-SEPT2* case DNA was available to perform methylation analysis, a definitive conclusion cannot be drawn.

How can *SEPT2* down-regulation be associated with leukemogenesis? Mammalian septins have been linked with two distinct steps in cell division, namely during chromosome segregation and during cytokinesis, as depletion of septins by siRNA result in defects in both of these processes [31,32]. *SEPT2* function is dependent of the formation of core oligomeric complexes with *SEPT6* and *SEPT7*, and this septin heterotrimer is a recognized regulator of microtubule stability, with septin depletion resulting in a marked stabilization of microtubules and mitotic defects *in vivo* [1,33]. It is known that proper organization of the cytoskeleton, including that of septin filaments, is required for cell-cycle progression and, as a consequence, septins are indirectly involved in driving or halting the cell-cycle engine [34]. In addition, the *SEPT2-SEPT6-SEPT7* complex can directly regulate cell-cycle progression by sequestering key signaling molecules involved in the DNA damage response and cell-cycle progression [33,34]. Down-regulation of the expression of septin genes has been described previously in neoplasia. Expression of the mitochondrial ARTS protein, a splice variant of the *SEPT4* gene, is lost in the majority of childhood acute lymphoblastic leukemias and *SEPT9* expression is down-regulated by promoter methylation in head and neck squamous cell carcinomas, suggesting that both genes can function as tumor suppressor genes [35,36]. *SEPT2* may also play a role in the pathogenesis of other AML subtypes in which it is not involved as fusion partner, since we have uncovered significant *SEPT2* RNA down-regulation in AML associated with the *PML-RARA* rearrangement and in AML with a normal karyotype.

SEPT2 is not the only septin family gene associated with hematological neoplasia. It has been shown that at least four other septins, *SEPT5*, *SEPT6*, *SEPT9* and *SEPT11*, are also *MLL* fusion partners [9-13]. There is increasing evidence supporting the hypothesis that *MLL* fusion partners are not randomly chosen, but rather functionally selected. For instance, the most frequent *MLL* fusion partners *AFF1* (*AF4*), *MLLT3* (*AF9*), *MLLT1* (*ENL*) and *MLLT10* (*AF10*) have been shown to belong to the same nuclear protein network [37]. Furthermore, the carboxyl-terminal domain of *ELL* and *MLLT10* have been shown to be required for the leukemic transformation associated with the *MLL-ELL* and *MLL-MLLT10* fusion proteins, respectively [38,39].

Conclusion

We provide evidence of *MLL* and *SEPT2* down-regulation in *MLL-SEPT2* myeloid neoplasia, as well as *MLL* under-expression in AML with *MLL* fusions with other partners other than *SEPT2* but, due to the small number of *MLL-SEPT2* cases available, these results should be confirmed in a larger series of patients. Characterization of the expression profile of other *MLL* fusion partners, including other septins, in hematological malignancies may allow a

better understanding of the pathobiological mechanisms of AML.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

NC designed and performed the research, analyzed the data and drafted the manuscript. JS performed the research and analyzed the data. SB performed the RT-PCR studies. VC performed the qMSP studies. FRR analyzed the data. SL, CC, LT, and JV performed the chromosome banding and molecular cytogenetic studies. SS, CHM, and AB characterized patients 60 and 61 and provide samples of both patients for RT-PCR and qRT-PCR studies. JMM and LN clinically assessed the patients. MRT coordinated the study and participated in manuscript writing. All authors read and approved the final manuscript.

Additional material

Additional file 1

Additional Table S1. Summary of clinical, molecular, and cytogenetic data of the 61 patients with hematological malignancies.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2407-9-147-S1.doc>]

Additional file 2

Additional Table S2. Oligonucleotide primers and probes (5'FAM, 3'TAMRA) used in this study.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2407-9-147-S2.doc>]

Acknowledgements

This research was supported by grants from the "Associação Portuguesa Contra a Leucemia" (2006-30.2.AP/MJ) and the "Comissão de Fomento da Investigação em Cuidados de Saúde" (P.I. 26/2007).

References

1. Weirich CS, Erzberger JP, Barral Y: **The septin family of GTPases: architecture and dynamics.** *Nat Rev Mol Cell Biol* 2008, **9**:478-489.
2. Gladfelter AS, Montagna C: **Seeking truth on Monte Verita. Workshop on the molecular biology and biochemistry of septins and septin function.** *EMBO Rep* 2007, **8**:1120-1126.
3. Joo E, Surka MC, Trimble WS: **Mammalian SEPT2 is required for scaffolding nonmuscle myosin II and its kinases.** *Dev Cell* 2007, **13**:677-690.
4. Hall PA, Russell SE: **The pathobiology of the septin gene family.** *J Pathol* 2004, **204**:489-505.
5. Cerveira N, Correia C, Bizarro S, Pinto C, Lisboa S, Mariz JM, Marques M, Teixeira MR: **SEPT2 is a new fusion partner of MLL in acute myeloid leukemia with t(2;11)(q37;q23).** *Oncogene* 2006, **25**:6147-6152.
6. van Binsbergen E, de Weerd O, Buijs A: **A new subtype of MLL-SEPT2 fusion transcript in therapy-related acute myeloid leukemia with t(2;11)(q37;q23): a case report and literature review.** *Cancer Genet Cytogenet* 2007, **176**:72-75.

7. Snijder S, Mellink CH, Lelie H van der: **Translocation (2;11)(q37;q23) in therapy-related myelodysplastic syndrome after treatment for acute promyelocytic leukemia.** *Cancer Genet Cytogenet* 2008, **180**:149-152.
8. Cerveira N, Santos J, Pinheiro M, Snijder S, Lelie H van der, Mellink C, Teixeira MR: **A novel MLL-SEPT2 fusion variant in therapy-related myelodysplastic syndrome.** *Cancer Genet Cytogenet* 2008, **185**:62-64.
9. Megonigal MD, Rappaport EF, Jones DH, Williams TM, Lovett BD, Kelly KM, Lerou PH, Moulton T, Budarf ML, Felix CA: **t(11;22)(q23;q11.2) in acute myeloid leukemia of infant twins fuses MLL with hCDCrel, a cell division cycle gene in the genomic region of deletion in DiGeorge and velocardiofacial syndromes.** *Proc Natl Acad Sci USA* 1998, **95**:6413-6418.
10. Osaka M, Rowley JD, Zeleznik-Le NJ: **MSF (MLL septin-like fusion), a fusion partner gene of MLL, in a therapy-related acute myeloid leukemia with a t(11;17)(q23;q25).** *Proc Natl Acad Sci USA* 1999, **96**:6428-6433.
11. Taki T, Ohnishi H, Shinohara K, Sako M, Bessho F, Yanagisawa M, Hayashi Y: **AF17q25, a putative septin family gene, fuses the MLL gene in acute myeloid leukemia with t(11;17)(q23;q25).** *Cancer Res* 1999, **59**:4261-4265.
12. Ono R, Taki T, Taketani T, Kawaguchi H, Taniwaki M, Okamura T, Kawa K, Hanada R, Kobayashi M, Hayashi Y: **SEPTIN6, a human homologue to mouse Septin6, is fused to MLL in infant acute myeloid leukemia with complex chromosomal abnormalities involving 11q23 and Xq24.** *Cancer Res* 2002, **62**:333-337.
13. Kojima K, Sakai I, Hasegawa A, Niiya H, Azuma T, Matsuo Y, Fujii N, Tanimoto M, Fujita S: **FLJ10849, a septin family gene, fuses MLL in a novel leukemia cell line CNLBC1 derived from chronic neutrophilic leukemia in transformation with t(4;11)(q21;q23).** *Leukemia* 2004, **18**:998-1005.
14. Shaffer LG, Tommerup N: *ISCN 2005 An International System for Human Cytogenetic Nomenclature* Basel: Karger; 1995.
15. van Dongen JJ, Macintyre EA, Gabert JA, Delabesse E, Rossi V, Saglio G, Gottardi E, Rambaldi A, Dotti G, Griesinger F, Parreira A, Gameiro P, Diaz MG, Malec M, Langerak AW, San Miguel JF, Biondi A: **Standardized RT-PCR analysis of fusion gene transcripts from chromosome aberrations in acute leukemia for detection of minimal residual disease. Report of the BIOMED-I Concerted Action: investigation of minimal residual disease in acute leukemia.** *Leukemia* 1999, **13**:1901-1928.
16. Chaplin T, Bernard O, Beverloo HB, Saha V, Hagemeyer A, Berger R, Young BD: **The t(10;11) translocation in acute myeloid leukemia (M5) consistently fuses the leucine zipper motif of AF10 onto the HRX gene.** *Blood* 1995, **86**:2073-2076.
17. Poirel H, Rack K, Delabesse E, Radford-Weiss I, Troussard X, Debort C, Leboeuf D, Bastard C, Picard F, Veil-Buzyn A, Flandrin G, Bernard O, Macintyre E: **Incidence and characterization of MLL gene (11q23) rearrangements in acute myeloid leukemia M1 and M5.** *Blood* 1996, **87**:2496-2505.
18. Busson-Le Coniat M, Salomon-Nguyen F, Hillion J, Bernard OA, Berger R: **MLL-AF1q fusion resulting from t(1;11) in acute leukemia.** *Leukemia* 1999, **13**:302-306.
19. Shih LY, Liang DC, Fu JF, Wu JH, Wang PN, Lin TL, Dunn P, Kuo MC, Tang TC, Lin TH, Lai CL: **Characterization of fusion partner genes in 114 patients with de novo acute myeloid leukemia and MLL rearrangement.** *Leukemia* 2006, **20**:218-223.
20. Gabert J, Beillard E, Velden VH van der, Bi W, Grimwade D, Pallisgaard N, Barbany G, Cazzaniga G, Cayuela JM, Cavé H, Pane F, Aerts JL, De Micheli D, Thirion X, Pradel V, González M, Viehmann S, Malec M, Saglio G, van Dongen JJ: **Standardization and quality control studies of 'real-time' quantitative reverse transcriptase polymerase chain reaction of fusion gene transcripts for residual disease detection in leukemia - a Europe Against Cancer program.** *Leukemia* 2003, **17**:2318-2357.
21. Beillard E, Pallisgaard N, Velden VH van der, Bi W, Dee R, Schoot E van der, Delabesse E, Macintyre E, Gottardi E, Saglio G, Watzinger F, Lion T, van Dongen JJ, Hokland P, Gabert J: **Evaluation of candidate control genes for diagnosis and residual disease detection in leukemic patients using 'real-time' quantitative reverse-transcriptase polymerase chain reaction (RQ-PCR) - a Europe against cancer program.** *Leukemia* 2003, **17**:2474-2486.
22. Clark SJ, Harrison J, Paul CL, Frommer M: **High sensitivity mapping of methylated cytosines.** *Nucleic Acids Res* 1994, **22**:2990-2997.
23. Takai D, Jones PA: **The CpG island searcher.** *In Silico Biology* 2003, **3**:21.
24. Prestridge DS: **Predicting Pol II promoter sequences using transcription factor binding sites.** *J Mol Biol* 1995, **249**:923-932.
25. Eads CA, Danenberg KD, Kawakami K, Saltz LB, Blake C, Shibata D, Danenberg PV, Laird PW: **MethylLight: a high-throughput assay to measure DNA methylation.** *Nucleic Acids Res* 2000, **28**:E32.
26. Eads CA, Lord RV, Wickramasinghe K, Long TI, Kurumboor Sk, Bernstein L, Peters JH, DeMeester SR, DeMeester TR, Skinner KA, Laird PW: **Epigenetic patterns in the progression of esophageal adenocarcinoma.** *Cancer Res* 2001, **61**:3410-3418.
27. Whitmann SP, Liu S, Vukosavljevic T, Rush LJ, Liu C, Klisovic MI, Maharry K, Guimond M, Strout MP, Becknell B, Dorrance A, Klisovic RB, Plass C, Bloomfield CD, Marcucci G, Caligiuri MA: **The MLL partial tandem duplication: evidence for recessive gain-of-function in acute myeloid leukemia identifies a novel patient subgroup for molecular-targeted therapy.** *Blood* 2005, **106**:345-352.
28. Schreiner SA, García-Cuellar MP, Fey GH, Slany RK: **The leukemogenic fusion of MLL with ENL creates a novel transcriptional transactivator.** *Leukemia* 1999, **13**:1525-1533.
29. Zeisig BB, Schreiner S, García-Cuellar MP, Slany RK: **Transcriptional activation is a key function encoded by MLL fusion partners.** *Leukemia* 2003, **17**:359-365.
30. Esteller M: **Epigenetics in cancer.** *N Engl J Med* 2008, **358**:1148-1159.
31. Kinoshita M, Kumar S, Mizoguchi A, Ide C, Kinoshita A, Haraguchi T, Hiraoka Y, Noda M: **Nedd5, a mammalian septin, is a novel cytoskeletal component interacting with actin-based structures.** *Genes Dev* 1997, **11**:1535-1547.
32. Spiliotis ET, Kinoshita M, Nelson WJ: **A mitotic septin scaffold required for mammalian chromosome congression and segregation.** *Science* 2005, **307**:1781-1785.
33. Kremer BE, Haystead T, Macara IG: **Mammalian septins regulate microtubule stability through interaction with the microtubule-binding protein MAP4.** *Mol Biol Cell* 2005, **16**:4648-4659.
34. Kremer BE, Haystead T, Macara IG: **Septins regulate actin organization and cell-cycle arrest through nuclear accumulation of NCK mediated by SOCS7.** *Cell* 2007, **130**:777-779.
35. Elhasid R, Sahar D, Merling A, Zivony Y, Rotem A, Ben-Arush M, Izraeli S, Bercovich D, Larisch S: **Mitochondrial pro-apoptotic ARTS protein is lost in the majority of acute lymphoblastic leukemia patients.** *Oncogene* 2004, **23**:5468-5475.
36. Bennett KL, Karpenko M, Lin MT, Claus R, Arab K, Dyckhoff G, Plinkert P, Herpel E, Smiraglia D, Plass C: **Frequently methylated tumor suppressor genes in head and neck squamous cell carcinoma.** *Cancer Res* 2008, **68**:4494-4499.
37. Meyer C, Schneider B, Jakob S, Strehl S, Attarbaschi A, Schnittger S, Schoch C, Jansen MWJC, van Dongen JJM, den Boer ML, Pieters R, Ennas M-G, Angelucci E, Koehl U, Greil J, Griesinger F, zur Stadt U, Eckert C, Szczepanski T, Niggli FK, Schäfer BVV, Kempfski H, Brady HJM, Zuna J, Trka J, Nigro LL, Biondi A, Delabesse E, Macintyre E, Stanulla M, Schrappe M, Haas OA, Burmeister T, Dingermann T, Klingebiel T, Marschalek R: **The MLL recombinome of acute leukemias.** *Leukemia* 2006, **20**:777-784.
38. DiMartino JF, Miller T, Ayton PM, Landewe T, Hess JL, Clearly ML, Shilatfard A: **A carboxy-terminal domain of ELL is required and sufficient for immortalization of myeloid progenitors by MLL-ELL.** *Blood* 2000, **96**:3887-3893.
39. DiMartino JF, Ayton PM, Chen EH, Naftzger CC, Young BD, Clearly ML: **The AF10 leucine zipper is required for leukemic transformation of myeloid progenitors by MLL-AF10.** *Blood* 2002, **99**:3780-3785.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2407/9/147/prepub>