

# Aromatase expression is increased in *BRCA1* mutation carriers

Ashwini L Chand\*<sup>1</sup>, kConFab<sup>2</sup>, Evan R Simpson<sup>1,3</sup> and Colin D Clyne<sup>1,3</sup>

Address: <sup>1</sup>Prince Henry's Institute of Medical Research, P.O. Box 5152, Clayton, Victoria 3168, Australia, <sup>2</sup>Kathleen Cuningham Foundation Consortium for Research into Familial Breast Cancer, Melbourne, Victoria, Australia and <sup>3</sup>Dept. of Biochemistry, Monash University, Clayton, Victoria, Australia

Email: Ashwini L Chand\* - ashwini.chand@princehenrys.org; kConFab - heather.thorne@petermac.org; Evan R Simpson - evan.simpson@princehenrys.org; Colin D Clyne - colin.clyne@princehenrys.org

\* Corresponding author

Published: 16 May 2009

Received: 11 December 2008

BMC Cancer 2009, 9:148 doi:10.1186/1471-2407-9-148

Accepted: 16 May 2009

This article is available from: <http://www.biomedcentral.com/1471-2407/9/148>

© 2009 Chand et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

**Background:** Until recently, the molecular mechanisms explaining increased incidence of ovarian and breast cancers in carriers of *BRCA1* gene mutations had not been clearly understood. Of significance is the finding that *BRCA1* negatively regulates aromatase expression *in vitro*. Our objective was to characterise aromatase gene (*CYP19A1*) and its promoter expression in breast adipose and ovarian tissue in *BRCA1* mutation carriers and unaffected controls.

**Methods:** We measured aromatase transcripts, total and promoter-specific (PII, PI.3, PI.4) in prophylactic oophorectomy or mastectomy, therapeutic mastectomy, ovarian and breast tissue from unaffected women.

**Results:** We demonstrate that the lack of functional *BRCA1* protein correlates to higher aromatase levels in 85% of *BRCA1* mutation carriers. This increase is mediated by aberrant transcriptional regulation of aromatase; in breast adipose by increases in promoter II/I.3 and I.4-specific transcripts; and in the ovary with elevation in promoter I.3 and II-specific transcripts.

**Conclusion:** Understanding the link between *BRCA1* and aromatase is significant in terms of understanding why carcinogenesis is restricted to estrogen-producing tissues in *BRCA1* mutation carriers.

## Background

The roles of *BRCA1* in cellular functions include cell cycle control, protein degradation, DNA damage repair, and transcriptional regulation of its target genes. One of its target genes is aromatase (*CYP19A1*), the enzyme that catalyses the conversion of  $C_{19}$  steroids into bioactive estrogens [1]. *In vitro* studies demonstrate the direct binding of *BRCA1* to the proximal promoter region of *CYP19A1* (promoter II) and as a consequence the repression of transcription [2,3]. Gene silencing of *BRCA1* leads to an ina-

bility to impair aromatase gene expression and enzyme activity [2-5]. However, whether this leads to aromatase excess in *BRCA1* mutation carriers is unknown.

This link between *BRCA1* and aromatase is significant in terms of understanding why carcinogenesis is restricted to estrogen-producing tissues in mutation carriers. Given that aromatase is critical in promoting tumour growth and *BRCA1* and 2 mutations account for an 80% increased risk in hereditary breast and ovarian cancer

development, it is important to investigate the relationship between BRCA1 and aromatase expression in patients.

The mechanism with which aromatase exerts its activity in a tissue-specific manner is via transcriptional regulation of multiple promoters on its gene [6]. In women, aromatase is expressed in ovarian granulosa cells (PII), placental syncytiotrophoblast (PI.1, and 2a) brain (PI.f), breast cancer (PII, PI.3), skin fibroblasts, bone osteoblasts and chondrocytes (PI.4) and adipose stromal fibroblasts (PI.4) [7,8].

The role of aromatase in promoting breast cancer is well defined; factors derived from malignant epithelial cells such as prostaglandin E<sub>2</sub> as well as trans-acting transcription factors such as Liver Receptor Homologue (LRH-1/NR5A2), cAMP response element binding protein (CREB), Activating Transcription Factor 2 (ATF2/CREB2) and CCAAT/enhancer binding protein  $\delta$  (C/EBP $\delta$ ) increase aromatase levels within the epithelial cells and surrounding adipose stromal fibroblasts [9-11]. Additionally in breast cancers, the tumour inhibits adipose stromal fibroblast differentiation while in normal breast tissues differentiation into mature adipocytes reduces aromatase expression [10,12].

Within the breast tissue, adipose stromal cells are the primary aromatase expressing cells and suppression of BRCA1 expression via siRNA results in up-regulation of aromatase mRNA [5]. Hu *et al* showed that this suppression of aromatase transcription while mediated by BRCA1-associated RING domain (BARD1) protein is also dependent on other tissue-specific co-regulators, present only in granulosa and adipose tissue and not cancer epithelial cells. In additions, BRCA1 displaces CBP/p300 from the transcriptional complex at promoter II [3].

The aims of the current study were to investigate whether women with hereditary BRCA1 mutations resulting in a reduction of BRCA1 protein levels or bioactivity, show alterations in *CYP19A1* gene expression in major aromatase target tissues such as breast and the ovary.

## Methods

### Breast and Ovarian Biopsy Samples

Samples of frozen tissue from breast and ovarian tissue biopsies were obtained from the Kathleen Cuningham Foundation Consortium for research into Familial Breast cancer (kConFab) tissue bank (Melbourne, Australia). Biopsies were obtained from women between the ages of 25–40 years who had undergone therapeutic or prophylactic mastectomy (n = 10 patients) or oophorectomy (n = 6 patients) due to being positive for known BRCA1 mutations.

Subjects had been screened for BRCA1 point mutations and BRCA1 Multiplex Ligation-dependent Probe Amplification (MLPA) for large genomic rearrangements. In all cases, prophylactic and therapeutic mastectomy or prophylactic oophorectomy biopsies were not derived from the cancer containing breast or ovary. This was to ensure that any changes in aromatase expression would not be caused by tumour-derived factors such as prostaglandin E<sub>2</sub> that are known to increase aromatase expression locally in adipose stromal cells surrounding breast cancers.

For the control cohort, breast adipose tissue samples were obtained from premenopausal women undergoing reduction mammoplasty. The age of patients from whom tissue was collected ranged from 23 to 49 years. Tissue was collected by Mr. A Kalus, The Avenue Plastic Surgery, Melbourne, Australia, snap frozen in liquid nitrogen and stored at -80°C until use. Tissue samples used for this study (n = 10) are part of a larger collection of control samples. This study was approved by the Southern Health Human Research and Ethics Committee (Monash Medical Centre).

### Reverse Transcription and Quantitative Real-Time PCR

Total RNA was isolated from tissue biopsy samples using the RNeasy Mini kit according to manufacturer's instructions (Qiagen). RNA preparations were DNase (Ambion) treated to eliminate any DNA contamination. First strand cDNA synthesis from 300 to 500 ng of total RNA was performed using avian myeloblastosis virus reverse transcriptase (Promega) primed by random hexamers according to manufacturer's instructions. Real-time PCR reactions were carried out using the following primer sets and annealing conditions outlined in Table 1.

Quantitative real-time PCR amplifications were performed on the Lightcycler (Roche) using Fast Start Master SYBR Green 1 (Roche) and specific primer pairs described above. As additional validation, quantitative real-time PCR was also performed using the ABI 7900 PCR machine (Applied Biosystems) using the SYBR chemistry (Applied Biosystems). Experiments run on both real-time PCR systems were with triplicate RT reactions that had been diluted 1 in 25. Experimental samples were quantified by comparison with purified standards of known concentration. All samples were normalised to 18S transcript levels.

### Statistical Analysis

Data points are shown as mean of triplicate determinations, n = 10/group for all parameters. Statistical comparisons were performed using GraphPad Prism software. All data were log transformed before analysis, and the variances for each group were analysed. Differences between control and treatment groups were analysed by Mann-Whitney U-test. Statistical significance was defined as  $P <$

**Table 1: Primer sequences and annealing temperatures**

Target Gene	Primer Sequence	Annealing temperature	Reference
<i>CYP19A1</i>	F: 5'-acccttctgcgtcgtgtca-3' R: 5'-tctgtggaaatcctgcgtctt-3'	54	[13,15]
<i>CYP19A1</i> Promoter I.3	F: 5'-gataagttctatcagacc-3' R: 5'-caggaatctgccgtgggaga-3'	53	[13,15]
<i>CYP19A1</i> Promoter I.4	F: 5'-gtgaccaactggagcctg-3' R: 5'-caggaatctgccgtgggaga-3'	55	[13,15]
<i>CYP19A1</i> Promoter II	F: 5'-gcaacaggagctatagat-3' R: 5'-caggaatctgccgtgggaga-3'	54	[13,15]
<i>18S</i>	F: 5'-cggctaccacatccaaggaa-3' R: 5'-gctggaattaccgcgct-3'	58	
<i>ER</i>	F: 5'-tgtccagccaccaaccagt-3' R: 5'-tttcaacattctccctcctctt-3'	55	
<i>LRH-I</i>	F: 5'-ctgatactggaacttttgaa-3' R: 5'-cttcatttggatcatcaacctt-3'	55	[10]
<i>CCND1</i>	F: 5'-aactacctggaccgcttct-3' R: 5'-ccacttgagcttggttcacca-3'	55	
<i>FSHR</i>	F: 5'-gcggaacccaacatcgtgtc-3' R: 5'-tgaagaaatctctgcgaaagt-3'	55	[14]

0.05. Spearman's rank correlation coefficient was used to analyse whether transcript levels derived from different primer specific qPCR were correlated.

## Results

### Promoter-specific expression of aromatase in breast adipose of BRCA1 mutations carriers

Breast adipose tissue was derived from prophylactic or therapeutic mastectomies from premenopausal women who had mutations in the *BRCA1* gene. qPCR analysis revealed 25.8-fold higher mean aromatase expression ( $1.29 \pm 0.5$ ,  $n = 10$  transcripts/18S,  $n = 10$ ) for prophylactic mastectomy samples compared to controls ( $0.05 \pm 0.01$ ,  $P < 0.0001$ , Figure 1a). Total aromatase expression level in therapeutic mastectomy samples was 650-fold higher compared to control samples ( $32.53 \pm 18.11$  and  $0.05 \pm 0.01$  transcripts/18S respectively,  $n = 10$ ,  $P < 0.019$ , Figure 1a). Aromatase transcript was uniformly detectable at low levels in normal premenopausal breast tissue. In contrast, in prophylactic and therapeutic mastectomy tissues, aromatase mRNA levels were increased in most samples. This is reflected in the high fold changes observed for aromatase and its promoter transcripts described below.

Increased aromatase expression in breast cancer-containing breast adipose is predominantly mediated by a switch in promoter usage from the constitutive adipose-specific promoter I.4-specific to gonadal-specific promoter II expression in the adipose stromal fibroblasts [15]. Consistent with this, in therapeutic mastectomy samples promoter II-specific expression was elevated 692-fold in all patients exhibiting increased total aromatase levels that was above control mean value ( $1654 \pm 921$  and  $2.39 \pm 0.68$  transcripts/18S respectively,  $n = 10$ ,  $P < 0.005$ , Figure 1b). We also observed 14-fold higher promoter II-specific


transcript expression in the *BRCA1* prophylactic mastectomy cohort ( $P < 0.011$ ) (Figure 1b). A significant positive correlation was observed between aromatase and promoter II-specific transcript levels in all subjects ( $r = 0.8$ ,  $P < 0.0001$ ).

Relative expression of promoter I.3-specific transcripts was 16-fold higher in prophylactic mastectomy samples compared to controls ( $0.0006 \pm 0.0002$  and  $0.01 \pm 0.0086$  transcripts/18S respectively,  $n = 9-10$ ,  $P < 0.008$ ). In therapeutic mastectomy samples, the mean expression ( $1.24 \pm 0.79$  transcripts/18S,  $n = 10$ ) was increased 2066-fold compared to control mean value ( $P < 0.008$ , Figure 1c). A significant and positive correlation was observed between aromatase and promoter I.3-specific transcript expression in all subjects ( $r = 0.46$ ,  $P = 0.01$ ).

Promoter I.4-specific expression was 34-fold higher in prophylactic mastectomy tissues compared to controls ( $1.36 \pm 0.77$  and  $0.04 \pm 0.03$  transcripts/18S respectively,  $n = 8-10$ ,  $P < 0.002$ , Figure 1d) while in the therapeutic mastectomy samples there was a 650 fold increase above basal levels observed (Figure 1d,  $P < 0.02$ ). Promoter I.4-specific transcripts were approximately 60-fold lower in abundance compared to promoter II-specific transcripts. There was no significant correlation between aromatase and promoter I.4 transcript expression ( $r = 0.34$ ,  $P = 0.07$ ).

### Expression of ER, cyclin D1 and LRH-I in BRCA1 mutations carriers and normal breast adipose

Analysis of the prophylactic and therapeutic mastectomy samples did not show significant differences in ER $\alpha$  levels compared to controls (Figure 2a). Cyclin D1 expression levels in prophylactic samples were significantly different to controls ( $0.03 \pm 0.02$  and  $0.0014 \pm 0.0004$ ,  $P < 0.04$ ).




**Figure 1**  
**Aromatase and its promoter usage in breast adipose of BRCA1 mutation carriers and control subjects.** Relative expression levels of (a) total aromatase transcripts (b) promoter II-specific transcripts (c) promoter I.3-specific transcripts and (d) promoter I.4-specific transcripts in breast adipose tissue samples from control women and prophylactic and therapeutic mastectomy samples from BRCA1 mutation carriers. Data has been normalized to 18S expression for each sample (n = 10 subjects per group, RT-PCR performed in triplicate for each sample). The mean expression level for each subject group is indicated with a horizontal line and \*P < 0.05, \*\*P < 0.01 and \*\*\*P < 0.0001 are significantly different versus control.

Likewise in therapeutic mastectomy samples, mean cyclin D1 expression were higher than controls ( $2.29 \pm 1.21$  and  $0.0015 \pm 0.0004$  transcripts/18S, n = 10,  $P < 0.008$ , Figure 2b). A positive correlation was observed with cyclin D1 and aromatase expression in all subjects ( $r = 0.48$ ,  $P = 0.007$ ).

Expression analysis of the orphan nuclear receptor LRH-1/NR5A2 did not demonstrate significant changes between prophylactic mastectomy and control subjects (Figure 2c). In therapeutic mastectomy samples the increase in expression compared to controls was 550-fold higher ( $0.0004 + 0.0001$  and  $0.22 + 0.20$  transcripts/18S, n = 10,  $P < 0.0002$ , Figure 2c). LRH-1/NR5A2 expression was also significantly and positively correlated with aromatase transcript levels ( $r = 0.53$ ,  $P = 0.002$ ).

**Transcriptional analysis of aromatase in the ovary of BRCA1 mutations carriers**

The mRNA transcripts specific for the gonad-specific promoter II, promoter I.3, and promoter I.4 were quantitated by real-time PCR analysis in control and prophylactic oophorectomy samples from BRCA1 mutation carriers. Increased CYP19A1 mRNA expression, promoter II- and I.3-specific transcripts were observed in ovarian tissue samples derived from BRCA1 mutation carriers compared to controls (Figures 3a–c). While aromatase transcript levels were not significantly different between control and BRCA1 groups (Figure 3a), promoter II-specific transcript levels were significantly increased in the prophylactic oophorectomy samples compared to controls levels ( $0.13 \pm 0.06$  and  $0.02 \pm 0.01$  transcripts/18S respectively,  $P < 0.05$ , n = 6, Figure 3b).



**Figure 2**

**ER $\alpha$ , cyclin D1 and LRH-1/NR5A2 expression in breast adipose in *BRCA1* mutation carriers and control subjects.** Relative expression levels of (a) ER $\alpha$ , (b) cyclin D1 and (c) LRH-1/NR5A2 in prophylactic and therapeutic mastectomy and control breast adipose tissue. Data has been normalized to 18S expression for each sample (n = 10 subjects per group, RT-PCR performed in triplicate for each sample). The mean expression level for each subject group is indicated with a horizontal line and \* $P < 0.05$ , \*\* $P < 0.01$  and \*\*\* $P < 0.0001$  are significantly different versus control.

Although approximately 100-fold lower in abundance than promoter II-specific transcripts, promoter I.3-specific transcript levels showed significantly higher expression in the *BRCA1* mutation carriers compared to controls ( $0.004 \pm 0.002$  and  $1.7 \times 10^{-5} \pm 3.8 \times 10^{-6}$  transcripts/18S respectively,  $P < 0.001$ , Figure 3c).

The primary site of estrogen production is in the ovarian follicles where the granulosa cells respond to the pituitary FSH stimulus to increase aromatase expression in a cyclic fashion [16-18]. As the study group consisted of premenopausal women, we also measured their FSHR transcript levels to assess whether the increase in aromatase expression observed was due to the FSHR status of the ovary. There was no significant difference in means of the two study groups (Figure 3d).

## Discussion


The aim of the current study was to investigate the clinical relationship between aromatase and increased risk of breast and ovarian cancer in *BRCA1* mutation carriers. We aimed to characterise aromatase transcriptional regulation in non-tumour containing breast adipose and ovary of women with pathogenic *BRCA1* mutations, who had undergone prophylactic or therapeutic mastectomy or prophylactic oophorectomy.

We show that aromatase expression is significantly higher in *BRCA1* mutation carriers, in patients who had experience breast cancer but also in women who had a high risk for breast cancer and had prophylactic removal of their breast tissue. As observed in studies with the tumour-associated breast adipose tissue, in the therapeutic mastecto-

mies, gonad-specific promoter I.3/II levels were significantly elevated. This supports the notion of a promoter-switch mechanism causing aromatase over-expression and increased local estrogen concentration as a tumour-promoting stimulus [6]. This finding also supports *in vitro* molecular study reports that show that *BRCA1* is part of a repression complex at promoter I.3/II [2-4].

As an extra validation of the inverse relationship between *BRCA1* and aromatase transcriptional activity, we also assessed breast and ovarian tissue derived from prophylactic organ removal from *BRCA1* mutation carriers. Increases in aromatase and its proximal promoter I.3/II transcripts were also observed in these *BRCA1* mutation carriers, supporting the hypothesis that with decreased *BRCA1* function there is a dysregulation of aromatase transcription regulation and a predisposing factor to breast and ovarian cancer.

In breast adipose from *BRCA1* mutation carriers, we observed greater fold increase in promoter I.3/II specific transcripts in therapeutic compared to prophylactic mastectomy tissue. The difference in the expression of transcripts derived from proximal promoters I.3 and II between the therapeutic and prophylactic samples suggests that there may be a difference in exposure to factors such as prostaglandin E<sub>2</sub> or a more downstream component of the signalling pathway such as cAMP between the two study groups. This may also indicate that in the presence of a tumour, increasing levels of factors such as prostaglandin E<sub>2</sub> also have effects on the adjacent, non-tumour breast adipose.



**Figure 3**  
**Aromatase, PII, PI.3 and FSHR expression in ovarian tissue of BRCA1 mutation carriers and control women.**  
 Transcript expression of (a) total aromatase, (b) promoter II-specific (c) promoter I.3-specific and (d) FSHR in tissue derived from prophylactic oophorectomy in BRCA1 mutation carriers and age-matched controls. Data has been normalized to 18S expression for each sample (n = 6 subjects per group, RT-PCR performed in triplicate for each sample). The mean expression level for each subject group is indicated with a horizontal line and \*P < 0.05, \*\*P < 0.001 are significantly different versus control.

Therefore the current transcriptional model for aromatase over-expression in tumour containing breast tissue may be similar for the tumour-free adjacent breast tissue. Tumour-derived prostaglandin E<sub>2</sub> stimulates the expression of orphan nuclear receptor LRH-1/NR5A2 and increases its occupancy on the nuclear receptor half-site upstream of promoter II [10]. This model is extended with data from *in vitro* experiments demonstrating that with prostaglandin E<sub>2</sub> treatment BRCA1 is removed from the histone acetylase p300 and phospho-CREB transcription complex that occupies the promoter I.3/II region [3].

The increase in LRH-1 expression in therapeutic mastectomy samples is further evidence of its role in driving pro-

motor II mediated aromatase transcription in adipose stromal cells surrounding breast tumours [10,19,20]. The promoter switch from the basal promoter I.4 to gonad-specific promoter II occurs with the increased expression of LRH-1 in adipose stromal cells as well as in the tumour epithelial cells [19-22]. Furthermore it has been shown that the LRH-1 gene is an estrogen-responsive [23].

The observed increase in basal breast-specific promoter I.4 transcript levels in both therapeutic and prophylactic mastectomy tissue samples implicates an inverse relationship between BRCA1 and promoter I.4 driven transcription. It could also imply that glucocorticoids and cytokines such as interleukin-6/11 and tumour necrosis factor-α which

stimulate promoter I.4-driven expression are elevated in *BRCA1* mutation carriers allowing the increase in promoter I.4-specific transcripts [24].

The mRNA expression of  $ER\alpha$  was investigated to understand whether increased hormone sensitivity was a mechanism for development of breast cancer in *BRCA1* mutation carriers. There were no changes observed in prophylactic mastectomy samples while in therapeutic mastectomies, some samples had increased  $ER\alpha$  levels, however this was not significant. In this case,  $ER\alpha$  functional studies would be a more relevant endpoint to address this hypothesis in light of previous studies that demonstrate that *BRCA1* interacts directly with  $ER\alpha$  suppressing  $ER\alpha$ -mediated transcription of target genes [25-27]. It was shown that the *BRCA1* protein binds to  $ER\alpha$  to inhibit activity of the activation function AF-2 domain and may cause conformational change and recruitment of coactivator proteins [27].

Cyclin D1 promotes progression through the G1-S phase of the cell cycle by phosphorylating and inactivating the retinoblastoma protein and its over-expression has been linked to early onset of cancer and increased risk of tumour progression and metastasis in parathyroid adenoma, breast cancer, colon cancer, lymphoma, melanoma, and prostate cancer (reviewed in [28]). The existing model for cyclin D1 function is via binding the cyclin dependent kinases, p300 and histone deacetylases to modulate local chromatin structure of its target genes that are involved in the regulation of cell proliferation and differentiation [29]. In the context of this study, cyclin D1 mRNA expression and its promoter activity have been shown to be up-regulated by adipokines produced by the adipose stromal cells leading to increased breast epithelial cell proliferation, motility and angiogenesis [30]. In addition, cyclin D1 interacts with  $ER\alpha$  and promotes its recruitment to estrogen response elements on promoters of target genes [31]. The significant increase in cyclin D1 expression in therapeutic mastectomy samples supports the idea that secreted factors in the breast stroma can promote increased cell proliferation and thus bring about a tumour-promoting environment. Also increased cyclin D1 levels may enhance  $ER\alpha$  activity especially in the presence of higher concentrations of estrogen.

In prophylactic oophorectomy tissues, the mean expression of total aromatase was 8-fold higher than control levels however this was not statistically significant. This may be due to the small sample size of the study group or that in premenopausal women aromatase is predominantly under the control of gonadotropins. There was a significant increase in promoter II- and I.3-specific transcripts indicating aberrant regulation of the transcriptional process. FSHR expression in both groups were not significantly

different (fold change difference 1.6 between groups). We conclude that the aromatase promoter transcript increases observed is not due to gonadotropin sensitivity.

In summary, the current study is the first that to validate the negative association between *BRCA1* function and aromatase expression in clinical samples. We demonstrate that the lack of functional *BRCA1* protein correlates to higher aromatase levels in 85% of *BRCA1* mutation carriers in our study cohort (therapeutic and prophylactic mastectomy tissues). We also show that the change in aromatase expression levels is mediated via aberrant transcriptional regulation of the *CYP19A1* gene; in breast adipose by increases in gonad and breast cancer-specific promoter II/I.3 and promoter I.4 transcripts; and in the ovary with elevation in breast cancer-specific promoter I.3 and promoter II transcripts.

We and others have shown positive correlation between quantitative RT-PCR, promoter-specific reporter assays, aromatase protein levels and aromatase enzymatic activity *in vitro* [10,19,24,32]. Furthermore little is known about post translational modification of the aromatase protein. Therefore the measurement of aromatase and its promoter-specific transcripts is considered an accurate reflection of aromatase activity *in vivo*.

The availability of pathology samples is a limitation for clinical studies in humans especially to perform experiments such as western blot analysis for protein expression. Despite the small sample size we have observed statistically significant differences between our control and study group highlighting that a larger scale study to address this question is of value. As data may be affected by other variables such as medication and parity, a more extensive patient history and a larger scale analysis is required to further understand the relationship between *BRCA1* and aromatase.

## Conclusion

Aromatase mRNA expression is increased in breast adipose tissue of *BRCA1* mutation carriers; this supports previous *in vitro* data showing interaction between *BRCA1* and the aromatase promoter. It also raises the possibility of prophylactic use of aromatase inhibitors as an alternative to surgical removal of tissue in high breast and ovarian cancer risk women.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

AC participated in the design of the study, carried out the experiments, performed data analysis and wrote the manuscript. CDC conceived the study, participated in its

design and manuscript preparation. ERS contributed intellectual input. kConFab recruited patients for the study and provided tissue biopsy samples. All authors read and approved the final manuscript.

### Acknowledgements

This research was supported by the National Health and Medical Research Council of Australia (NHMRC) (CDC, 338518). We wish to thank Prof. Peter Fuller for normal ovarian biopsy tissue from premenopausal women. Additionally we thank Heather Thorne, Eveline Niedermayr, all the kConFab research nurses and staff, the heads and staff of the Family Cancer Clinics, and the Clinical Follow Up Study (funded by NHMRC grants 145684, 288704 and 454508) for their contributions to this resource, and the many families who contribute to kConFab. kConFab is supported by grants from the National Breast Cancer Foundation, NHMRC and by the Queensland Cancer Fund, the Cancer Councils of New South Wales, Victoria, Tasmania and South Australia, and the Cancer Foundation of Western Australia. We are also grateful to Mr. A Kalus, Avenue Plastic Surgery, Melbourne and patients for normal breast tissue samples.

### References

- Simpson ER, Dowsett M: **Aromatase and its inhibitors: significance for breast cancer therapy.** *Recent Prog Horm Res* 2002, **57**:317-338.
- Lu M, Chen D, Lin Z, Reierstad S, Trauernicht AM, Boyer TG, Bulun SE: **BRCA1 negatively regulates the cancer-associated aromatase promoters I.3 and II in breast adipose fibroblasts and malignant epithelial cells.** *J Clin Endocrinol Metab* 2006, **91**:4514-4519.
- Subbaramaiah K, Hudis C, Chang SH, Hla T, Dannenberg AJ: **EP2 and EP4 receptors regulate aromatase expression in human adipocytes and breast cancer cells. Evidence of a BRCA1 and p300 exchange.** *J Biol Chem* 2008, **283**:3433-3444.
- Ghosh S, Lu Y, Katz A, Hu Y, Li R: **Tumor suppressor BRCA1 inhibits a breast cancer-associated promoter of the aromatase gene (CYP19) in human adipose stromal cells.** *Am J Physiol Endocrinol Metab* 2007, **292**:246-252.
- Hu Y, Ghosh S, Amlah A, Yue W, Lu Y, Katz A, Li R: **Modulation of aromatase expression by BRCA1: a possible link to tissue-specific tumor suppression.** *Oncogene* 2005, **24**:8343-8348.
- Simpson ER, Davis SR: **Minireview: aromatase and the regulation of estrogen biosynthesis – some new perspectives.** *Endocrinology* 2001, **142**:4589-4594.
- Bulun SE, Simpson ER: **Regulation of aromatase expression in human tissues.** *Breast Cancer Res Treat* 1994, **30**:19-29.
- Simpson ER, Mahendroo MS, Means GD, Kilgore MW, Corbin CJ, Mendelson CR: **Tissue-specific promoters regulate aromatase cytochrome P450 expression.** *J Steroid Biochem Mol Biol* 1993, **44**:321-330.
- Chen D, Reierstad S, Lin Z, Lu M, Brooks C, Li N, Innes J, Bulun SE: **Prostaglandin E(2) induces breast cancer related aromatase promoters via activation of p38 and c-Jun NH(2)-terminal kinase in adipose fibroblasts.** *Cancer Res* 2007, **67**:8914-8922.
- Clyne CD, Speed CJ, Zhou J, Simpson ER: **Liver receptor homologue-I (LRH-I) regulates expression of aromatase in preadipocytes.** *J Biol Chem* 2002, **277**:20591-20597.
- Kijima I, Ye J, Glackin C, Chen S: **CCAAT/enhancer binding protein delta up-regulates aromatase promoters I.3/II in breast cancer epithelial cells.** *Cancer Res* 2008, **68**:4455-4464.
- Meng L, Zhou J, Sasano H, Suzuki T, Zeitoun KM, Bulun SE: **Tumor necrosis factor alpha and interleukin 11 secreted by malignant breast epithelial cells inhibit adipocyte differentiation by selectively down-regulating CCAAT/enhancer binding protein alpha and peroxisome proliferator-activated receptor gamma: mechanism of desmoplastic reaction.** *Cancer Res* 2001, **61**:2250-2255.
- Agarwal VR, Bulun SE, Simpson ER: **Quantitative detection of alternatively spliced transcripts of the aromatase cytochrome P450 (CYP19) gene in aromatase-expressing human cells by competitive RT-PCR.** *Mol Cell Probes* 1999, **9**:453-464.
- Fuller PJ, Verity K, Shen Y, Mamers P, Jobling T, Burger HG: **No evidence of a role for mutations or polymorphisms of the follicle-stimulating hormone receptor in ovarian granulosa cell tumors.** *J Clin Endocrinol Metab* 1998, **83**:274-279.
- Mahendroo MS, Mendelson CR, Simpson ER: **Tissue-specific and hormonally controlled alternative promoters regulate aromatase cytochrome P450 gene expression in human adipose tissue.** *J Biol Chem* 1993, **268**:19463-19470.
- Kwintkiewicz J, Cai Z, Stocco C: **Follicle-stimulating hormone-induced activation of Gata4 contributes in the up-regulation of Cyp19 expression in rat granulosa cells.** *Mol Endocrinol* 2007, **21**:933-947.
- Parakh TN, Hernandez JA, Grammer JC, Weck J, Hunzicker-Dunn M, Zeleznik AJ, Nilson JH: **Follicle-stimulating hormone/cAMP regulation of aromatase gene expression requires beta-catenin.** *Proc Natl Acad Sci USA* 2006, **103**:12435-12440.
- Richards JS: **Perspective: the ovarian follicle – a perspective in 2001.** *Endocrinology* 2001, **142**:2184-2193.
- Clyne CD, Kovacic A, Speed CJ, Zhou J, Pezzi V, Simpson ER: **Regulation of aromatase expression by the nuclear receptor LRH-I in adipose tissue.** *Mol Cell Endocrinol* 2004, **215**:39-44.
- Zhou J, Suzuki T, Kovacic A, Saito R, Miki Y, Ishida T, Moriya T, Simpson ER, Sasano H, Clyne CD: **Interactions between prostaglandin E(2), liver receptor homologue-1, and aromatase in breast cancer.** *Cancer Res* 2005, **65**:657-663.
- Miki Y, Clyne CD, Suzuki T, Moriya T, Shibuya R, Nakamura Y, Ishida T, Yabuki N, Kitada K, Hayashi S, Sasano H: **Immunolocalization of liver receptor homologue-1 (LRH-I) in human breast carcinoma: possible regulator of in situ steroidogenesis.** *Cancer Lett* 2006, **244**:24-33.
- Bouchard MF, Taniguchi H, Viger RS: **Protein kinase A-dependent synergism between GATA factors and the nuclear receptor, liver receptor homolog-1, regulates human aromatase (CYP19) PII promoter activity in breast cancer cells.** *Endocrinology* 2005, **146**:4905-4916.
- Annicotte JS, Chavey C, Servant N, Teyssier J, Bardin A, Licznar A, Badia E, Pujol P, Vignon F, Maudelonde T, Lazennec G, Cavailles V, Fajas L: **The nuclear receptor liver receptor homolog-1 is an estrogen receptor target gene.** *Oncogene* 2005, **24**:8167-8175.
- Chen D, Reierstad S, Lu M, Lin Z, Ishikawa H, Bulun SE: **Regulation of breast cancer-associated aromatase promoters.** *Cancer Lett* 2009, **273**:15-27.
- Fan S, Ma YX, Wang C, Yuan RQ, Meng Q, Wang JA, Erdos M, Goldberg ID, Webb P, Kushner PJ, Pestell RG, Rosen EM: **Role of direct interaction in BRCA1 inhibition of estrogen receptor activity.** *Oncogene* 2001, **20**:77-87.
- Rosen EM, Fan S, Isaacs C: **BRCA1 in hormonal carcinogenesis: basic and clinical research.** *Endocr Relat Cancer* 2005, **12**:533-548.
- Fan S, Wang J, Yuan R, Ma Y, Meng Q, Erdos MR, Pestell RG, Yuan F, Auburn KJ, Goldberg ID, Rosen EM: **BRCA1 inhibition of estrogen receptor signaling in transfected cells.** *Science* 1999, **284**:1354-1356.
- Fu M, Wang C, Li Z, Sakamaki T, Pestell RG: **Minireview: Cyclin D1: normal and abnormal functions.** *Endocrinology* 2004, **145**:5439-5447.
- Reutens AT, Fu M, Wang C, Albanese C, McPhaul MJ, Sun Z, Balk SP, Janne OA, Palvimo JJ, Pestell RG: **Cyclin D1 binds the androgen receptor and regulates hormone-dependent signaling in a p300/CBP-associated factor (P/CAF)-dependent manner.** *Mol Endocrinol* 2001, **15**:797-811.
- Iyengar P, Combs TP, Shah SJ, Gouon-Evans V, Pollard JW, Albanese C, Flanagan L, Tenniswood MP, Guha C, Lisanti MP, Pestell RG, Scherer PE: **Adipocyte-secreted factors synergistically promote mammary tumorigenesis through induction of anti-apoptotic transcriptional programs and proto-oncogene stabilization.** *Oncogene* 2003, **22**:6408-6423.
- Zwijsen RM, Buckle RS, Hijmans EM, Loomans CJ, Bernards R: **Ligand-independent recruitment of steroid receptor coactivators to estrogen receptor by cyclin D1.** *Genes Dev* 1998, **12**:3488-3498.
- McInnes KJ, Brown KA, Knower KC, Chand AL, Clyne CD, Simpson ER: **Characterisation of aromatase expression in the human adipocyte cell line SGBS.** *Breast Cancer Res Treat* 2008.



### Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2407/9/148/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

