

STUDY PROTOCOL

Open Access



Study protocol of HGCSG1404 SNOW study: a phase I/II trial of combined chemotherapy of S-1, nab-paclitaxel and oxaliplatin administered biweekly to patients with advanced gastric cancer

Yasuyuki Kawamoto^{1,2}, Yoshito Komatsu^{1*}, Satoshi Yuki², Kentaro Sawada², Tetsuhito Muranaka^{1,2}, Kazuaki Harada^{1,2}, Hiroshi Nakatsumi^{1,2}, Hiraku Fukushima³, Atsushi Ishiguro⁴, Masayoshi Dazai⁵, Kazuteru Hatanaka⁶, Michio Nakamura⁷, Ichiro Iwanaga⁸, Minoru Uebayashi⁸, Susumu Sogabe⁹, Yoshimitsu Kobayashi⁹, Takuto Miyagishima⁹, Kota Ono¹⁰, Naoya Sakamoto² and Yuh Sakata¹¹

Abstract

Background: In Japan, S-1 plus cisplatin (SP) regimen has become a standard therapy for patients with advanced gastric cancer. Moreover, the S-1 plus oxaliplatin regimen is now a standard treatment.

Nab-paclitaxel was developed for chemotherapy of gastric cancer in Japanese clinical practice. Nab-paclitaxel, created with albumin-bound paclitaxel particles, has high transferability to tumour tissues and does not cause hypersensitivity reactions because of a different chemical composition compared with docetaxel and paclitaxel. A combination of S-1, nab-paclitaxel and oxaliplatin (which we named "SNOW regimen") can be a promising triplet therapy for advanced gastric cancer. Although we have to pay attention to chemotherapy-induced neuropathy, we aim to investigate the recommended dose of this regimen in a phase I study. Furthermore, we will investigate its efficacy and toxicity in a phase II study.

Methods: The phase I study is a dose-escalation study using a standard 3 plus 3 design, followed by expansion cohorts. The SNOW regimen involves 28-day cycles with escalated doses of nab-paclitaxel (100–175 mg/m² on days 1 and 15) and fixed doses of oxaliplatin (65 mg/m² on days 1 and 15) and S-1 (80 mg/m²/day on day 1 to 14). The primary endpoints are assessment of dose limiting toxicities and determination of maximum tolerated dose to investigate the recommended dose in the subsequent phase II study. In the phase II study, the primary endpoint is objective response rate. Secondary endpoints are assessment of safety, progression-free survival, disease control rate, overall survival and time to treatment failure. Adverse events were monitored and graded according to the National Cancer Institute Common Terminology Criteria for Adverse Events version 4.0.

Discussion: Triplet therapies for advanced gastric cancer patients have been evaluated in clinical trials. The SNOW regimen can be a promising new triplet therapy.

Trial registration: This study is performed at institutes that participate in Hokkaido Gastrointestinal Cancer Study Group (HGCSG) and registered as UMIN000016788. Registered 16 March 2015.

Keywords: Gastric cancer, Chemotherapy, S-1, Nab-paclitaxel, Oxaliplatin

* Correspondence: ykomatsu@ac.cyberhome.ne.jp

¹Department of Cancer Center, Hokkaido University Hospital, Sapporo, Japan
Full list of author information is available at the end of the article

Background

In 2012, gastric cancer was the third leading cause of cancer deaths worldwide, responsible for 723,000 deaths [1]. Approximately 60% of gastric cancer patients worldwide are diagnosed in East Asian countries (Japan, China and Korea) [2].

Standard cytotoxic chemotherapy is frequently used as a first-line treatment for advanced gastric cancer (AGC), with a median overall survival of 8–12 months. The survival benefit with chemotherapy is not yet adequate; therefore, new agents and combination therapies are needed to improve the outcome of patients with advanced disease.

In patients with human epidermal growth factor receptor type 2 (HER2)-positive AGC, trastuzumab demonstrated a survival benefit in the ToGA study [3]. Trastuzumab in combination with cisplatin plus capecitabine or fluorouracil is a worldwide standard treatment for HER2-positive AGC. Meanwhile, for AGC without HER2 overexpression, several doublet or triplet first-line chemotherapy regimens, including fluorouracil, platinum, anthracycline or taxanes, are available [4, 5]. However, especially in triplet regimens, toxicity profiles must be carefully considered.

S-1 is an oral anticancer drug that combines tegafur, a prodrug of fluorouracil, with 5-chloro-2,4-dihydropyrimidine (CDHP) and oteracil potassium in a molar ratio of 1:0.4:1. CDHP reversibly antagonizes the activity of dihydropyrimidine dehydrogenase, the rate-limiting enzyme for the degradation of fluorouracil [6].

In Japan, the S-1 plus cisplatin (SP) regimen has become a standard therapy for patients with AGC [7]. Moreover, the S-1 combined with oxaliplatin (SOX) regimen has become a standard treatment [8].

Nab-paclitaxel was also developed for chemotherapy of gastric cancer in Japanese clinical practice [9]. Nab-paclitaxel, which is created with albumin-bound paclitaxel particles, has high transferability to tumour tissues and hardly cause a hypersensitivity reaction because of different chemical composition compared with docetaxel and paclitaxel.

To further improve the antitumor efficacy, the DCS regimen (SP combined with docetaxel) is considered as one of promising candidate of new standard treatment. And we have devised an S-1, nab-paclitaxel and oxaliplatin combination (which we named the ‘SNOW regimen’) which we have changed docetaxel to nab-paclitaxel, and cisplatin to oxaliplatin. That could be a promising triplet therapy for AGC patients. Although we have to pay attention to chemotherapy-induced neuropathy, we are aiming to investigate the recommended dose of this regimen in a phase I study. Furthermore, we will consider the efficacy and toxicity of this regimen in a phase II study.

Methods/Design

Inclusion criteria

1) Unresectable advanced, metastatic, or recurrent gastric or gastroesophageal junction cancer that is pathologically diagnosed as adenocarcinoma.

2) HER2 negative [0, 1+ in immunohistochemistry (IHC) or 2+ in IHC and FISH negative].

3) Patients who have measurable lesions based on Response Evaluation Criteria in Solid Tumors (RECIST) version 1.1.

4) Patients who have received no prior chemotherapy or radiotherapy for gastric or gastroesophageal junction cancer (patients who have received adjuvant therapy including S-1 before 180 days or more can be eligible; however, patients who have received adjuvant therapy including oxaliplatin at any time cannot be eligible).

5) 20 years of age and older.

6) Patients with Eastern Cooperative Oncology Group Performance Status of 0 or 1.

7) Patients who have possibility of oral intake.

8) Patients must have sufficient organ function as below:

- Absolute neutrophil count $\geq 1500/\text{mm}^3$
- Platelet count $\geq 100,000/\text{mm}^3$
- Haemoglobin ≥ 9.0 g/dL
- Total bilirubin ≤ 1.5 mg/dL
- Aspartate aminotransferase ≤ 100 U/L (≤ 200 in patients with liver metastases)
- Alanine aminotransferase ≤ 100 U/L (≤ 200 in patients with liver metastases)
- Serum creatinine ≤ 1.2 mg/dL
- Creatinine clearance ≥ 60 mL/min

9) Patients with a life expectancy of at least three months.

10) Patients must provide written informed consent.

Exclusion criteria

1) Patients with history of hypersensitivity to any drugs in this study.

2) Patients with active infection.

3) Patients who are hepatitis B antigen positive.

4) Patients with serious complications, such as

- Uncontrollable cardiovascular disease, angina and arrhythmia
- Myocardial infarction in past three months
- Uncontrollable diabetes mellitus
- Intestinal lung disease or pulmonary fibrosis

5) Patients with \geq Grade 2 peripheral neuropathy.

- 6) Patients with any central nervous system metastases that are symptomatic or required treatment.
- 7) Patients with uncontrollable diarrhoea.
- 8) Patients with multiple primary cancers.
- 9) Female patients who are pregnant or lactating, or planning to become pregnant or lactating.
- 10) Other patients who are considered to be unsuitable for this study by the investigator.

Treatment

The SNOW regimen consists of 28-day cycles with escalated doses of nab-paclitaxel (100–175 mg/m² on days 1 and 15) and fixed doses of oxaliplatin (65 mg/m² on days 1 and 15) and S-1 (80 mg/m²/day on day 1 to 14) (Fig. 1). In the setting of the administration schedule of the triplet regimen, the point where given in divided doses might relieve chemotherapy-induced neuropathy than both drugs gave a high dose in once, because nab-paclitaxel had an adverse event of chemotherapy-induced neuropathy and oxaliplatin coming at the same time was considered. The administration every two weeks of oxaliplatin was frequently used for FOLFOX therapies for colorectal cancer. And there was a report used by the administration method with 175 mg/m² and 220 mg/m² as preoperative chemotherapy as multidrug therapy for breast cancer for every two weeks about nab-paclitaxel. For these reason, we decided to set a dose in this schedule. For gastric cancer, oxaliplatin is approved in 130 mg/m² every 3 weeks, and the dose concerned is 43.3 mg/m² a week. Because this study was combination therapy, and chemotherapy-induced neuropathy might strongly develop by combination with nab-paclitaxel, we set 65 mg/m² every 2 weeks at a fixation dose with a dose of oxaliplatin.

Phase I part

The phase I part of the study is a dose-escalation study using a standard 3 plus 3 design followed by expansion cohorts as below (Figs. 2 and 3).

- We start at level 1.
- The recommended dose (RD) is defined as one dose level lower than the maximum tolerated dose (MTD).
- If 1 of three patients experienced dose-limiting toxicities (DLT), three more patients were enrolled at the same dose level.
- The MTD is defined as the dose level at which two or more of three patients, or at least two of 4–6 patients, had DLTs during cycle 1.

Adverse events are monitored and graded according to NCI-CTCAE (the National Cancer Institute Common Terminology Criteria for Adverse Events) version 4.0. The following adverse drug reactions are defined as DLT:

- ≥Grade 3 febrile neutropenia
- Grade 4 thrombocytopenia
- Grade 4 neutropenia over 7 days
- ≥Grade 3 non-haematological toxicities (excludes nausea and vomiting)
- delay of starting cycle 2 longer than 15 days due to adverse event

The primary endpoints are assessment of DLTs and determination of MTD to investigate the RD in subsequent phase II study.

In the phase I study, we also investigate drug concentrations in all patients (Fig. 4).

Phase II part

In the phase II study, the primary endpoint is objective response rate according to RECIST version 1.1. We define the ratio of patients who are complete response or partial response in the overall study treatment time. Secondary endpoints are the assessment of adverse events graded according to NCI-CTCAE version 4.0, disease

SNOW: S-1 + Nab-paclitaxel + Oxaliplatin



1. Nab-paclitaxel: 100–175 mg/m² on days 1 and 15
2. Oxaliplatin: 65 mg/m² on days 1 and 15
3. S-1: 40–60 mg/body bid between day 1 and 15

Fig. 1 SNOW regimen. SNOW regimen comprises 28-day cycles with nab-paclitaxel and oxaliplatin on days 1 and 15, and S-1 (80 mg/m²/day on days 1–14)

【S-1 dose】

Body surface area	S-1
1.25 m ² <	40 mg bid/day [80 mg/day]
1.25m ² ≥, 1.5m ² <	50 mg bid/day [100 mg/day]
1.5m ² ≥	60 mg bid/day [120 mg/day]

【Dose level】

Level	Nab-PTX	L-OHP	S-1
0	100 mg/m ²	65 mg/m ²	80 mg/m ²
1	125 mg/m ²	65 mg/m ²	80 mg/m ²
2	150 mg/m ²	65 mg/m ²	80 mg/m ²
3	175 mg/m ²	65 mg/m ²	80 mg/m ²

Fig. 2 Dose-escalation in phase I part of SNOW regimen. In part of phase I study, escalated dose of nab-paclitaxel (100–175 mg/m² on days 1 and 15), fixed dose of oxaliplatin (65 mg/ m² on days 1 and 15) and S-1 (80 mg/m²/day on days 1–14) are administered

control rate, progression-free survival (PFS), time to treatment failure (TTF) and overall survival (OS). We evaluate PFS, TTF and OS using the Kaplan–Meier method.


Estimated number of enrollments

In phase I, 3 to 6 patients are enrolled at each dose level (maximum of 18 patients). In phase II, we referred to the SOX [8] and DCS [10] regimens. The response rate was 58.8% (95% CI; 44.2–72.4%) for SOX and 81.4% (95% CI; 69.1–90.3%) for DSC. With a threshold response rate of 59% and an expected

response rate of 81%, the simulation results indicated a sample size of 45 with α = 0.05 (both sides) for a power of 90% based on One Arm Binomial of SWOG. With an estimated dropout of some cases, a target sample size of 50 was estimated. That includes the cases administered the RD in phase I.

Discussion

Triplet therapies for AGC patients have been evaluated in clinical trials [4, 5, 10–12]. The SNOW regimen comprising an S-1, nab-paclitaxel and oxaliplatin combination could be a promising triplet therapy.



DLT	/ patients	
0	/ 3	to next level
1	/ 3	three more patients are enrolled
2 or more	/ 3	MTD
1	/ 6	to next level
2 or more	/ 4-6	MTD

Fig. 3 Flow chart in phase I part of SNOW regimen. In part of phase I study, this is a dose-escalation study using a standard 3 plus 3 design followed by expansion cohorts

In the phase I study, we investigated drug concentration in all patients.

- Nab-PTX: Paclitaxel
- L-OHP: Total-platinum, Free-platinum
- S-1: 5-FU




Fig. 4 Pharmacodynamics study in phase I part of SNOW regimen. In the phase I study, we also investigated drug concentration in all patients. We measured paclitaxel concentration from nab-paclitaxel, total-platinum and free-platinum from L-OHP and 5-FU from S-1

Oxaliplatin often causes neuronopathy-type peripheral sensory neuropathy. Meanwhile, nab-paclitaxel causes axonopathy-type peripheral sensory neuropathy. The combination of oxaliplatin and nab-paclitaxel might cause severe chemotherapy-induced peripheral neuropathy; therefore, we have to be vigilant for signs of neuropathy. Administration of nab-paclitaxel in divided doses might reduce the neuro-toxicity.

This study is performed at institutes that participate in the Hokkaido Gastrointestinal Cancer Study Group (HGCSG) and is registered as UMIN000016788. Registered 16 March 2015.

Abbreviations

AGC: Advanced gastric cancer; CDHP: 5-chloro-2,4-dihydropyrimidine; DLT: Dose-limiting toxicities; HER2: Human epidermal growth factor receptor type 2; IHC: Immunohistochemistry; MTD: Maximum tolerated dose; OS: Overall survival; PFS: Progression-free survival; RD: Recommended dose; TTF: Time to treatment failure

Acknowledgements

We are grateful to all of the patients and the co-investigators for their cooperation in HGCSG1404 SNOW study. The authors would like to thank Enago (www.enago.jp) for the English language review.

Funding

HGCSG1404 SNOW study is conducted as a research of Hokkaido Gastrointestinal Cancer Study Group with funding from Taiho Pharmaceutical CO., LTD. under the study contract. There is no competing interest between this company and the investigators that require disclosure in connection with the study.

Availability of data and materials

Not applicable.

Authors' contributions

YK, as a task manager, participated in entire coordinating of the study, design and writing of the protocol, data collection, data analysis, data interpretation, and writing of the manuscript. YK, SY, KS, TM, KH, HN, HF, AI, MD, KH, MN, II, MU, SS, TM and KO, as protocol preparation committee, participated in all phases of this study, including design and writing of the protocol, data collection, data analysis, data interpretation, and preparation of the manuscript. KO, as a chief of statistical analysis, participated in statistical setting of study design and data analysis. All authors reviewed and approved the final manuscript.

Ethics approval and consent to participate

This study is conducted in accordance with the Declaration of Helsinki and Ethical Guidelines for Medical and Health Research Involving Human Subjects and has been approved by the Institutional Review Boards of each participating institute. HGCSG1404 received approval on 14/Sep/2014 by Hokkaido University Hospital Research Ethics Committee. All patients provided written informed consent before enrollment.

Consent for publication

Not applicable.

Competing interests

YK reports personal fees from Taiho, Yakult, Bristol-Myers, Merck, Chugai, Takeda, Novartis, Pfizer, Bayer, and Daiichi-Sankyo. MN YS.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Department of Cancer Center, Hokkaido University Hospital, Sapporo, Japan. ²Department of Gastroenterology and Hepatology, Hokkaido University Graduate School of Medicine, Sapporo, Japan. ³Department of Gastroenterology, JCHO, Sapporo Hokushin Hospital, Sapporo, Japan. ⁴Department of Medical Oncology, Teine Keijinkai Hospital, Sapporo, Japan. ⁵Department of Gastroenterology, Sapporo Medical Center NTT EC, Sapporo, Japan. ⁶Department of Gastroenterology, Hakodate Municipal Hospital, Hakodate, Japan. ⁷Department of Gastroenterology, Sapporo City General Hospital, Sapporo, Japan. ⁸Department of Gastroenterology, Japanese Red Cross Kitami Hospital, Kitami, Japan. ⁹Department of Internal Medicine, Kushiro Rosai Hospital, Kushiro, Japan. ¹⁰Hokkaido University Hospital Clinical Research and Medical Innovation Center, Sapporo, Japan. ¹¹CEO, Misawa City Hospital, Misawa, Japan.

Received: 3 October 2016 Accepted: 27 November 2017

Published online: 08 December 2017

References

1. Cancer Genome Atlas Research Network. Comprehensive molecular characterization of gastric adenocarcinoma. *Nature*. 2014;513:202–9.
2. Ferlay J, Parkin DM, Steliarova-Foucher E. Estimates of cancer incidence and mortality in Europe in 2008. *Eur J Cancer*. 2010;46:765–81.
3. Bang YJ. Trastuzumab in combination with chemotherapy versus chemotherapy alone for treatment of HER2-positive advanced gastric or gastro-oesophageal cancer (ToGA): a phase 3, open-label, randomised controlled trial. *Lancet*. 2010; 376:687–97.
4. Van Cutsem E, Moiseyenko VM, Tjulandin S, Majlis A, Constenla M, Boni C, et al. Phase III study of docetaxel and cisplatin plus fluorouracil compared

- with cisplatin and fluorouracil as first-line therapy for advanced gastric cancer: a report of the V325 study group. *J Clin Oncol.* 2006;24:4991–7.
5. Cunningham D, Starling N, Rao S, Iveson T, Nicolson M, Coxon F, et al. Capecitabine and oxaliplatin for advanced esophagogastric cancer. *N Engl J Med.* 2008;358:36–46.
 6. Shirasaka T, Shimamoto Y, Ohshimo H, Yamaguchi M, Kato T, Yonekura K, et al. Development of a novel form of an oral 5-fluorouracil derivative (S-1) directed to the potentiation of the tumor selective cytotoxicity of 5-fluorouracil by two biochemical modulators. *Anti-Cancer Drugs.* 1996;7:548–57.
 7. Koizumi W, Narahara H, Hara T, Takagane A, Akiya T, Takagi M, et al. S-1 plus cisplatin versus S-1 alone for first-line treatment of advanced gastric cancer (SPIRITS trial): a phase III trial. *Lancet Oncol.* 2008;9:215–21.
 8. Yamada Y, Higuchi K, Nishikawa K, Gotoh M, Fuse N, Sugimoto N, et al. Phase III study comparing oxaliplatin plus S-1 with cisplatin plus S-1 in chemotherapy-naïve patients with advanced gastric cancer. *Ann Oncol.* 2015;26:141–8.
 9. Sasaki Y, Nishina T, Yasui H, Goto M, Muro K, Tsuji A, et al. Phase II trial of nanoparticle albumin-bound paclitaxel as second-line chemotherapy for unresectable or recurrent gastric cancer. *Cancer Sci.* 2014;105:812–7.
 10. Koizumi W, Nakayama N, Tanabe S, Sasaki T, Higuchi K, Nishimura K, et al. A multicenter phase II study of combined chemotherapy with docetaxel, cisplatin, and S-1 in patients with unresectable or recurrent gastric cancer (KDOG 0601). *Cancer Chemother Pharmacol.* 2012;69:407–13.
 11. Shah MA, Janjigian YY, Stoller R, Shibata S, Kemeny M, Krishnamurthi S, et al. Randomized multicenter phase II study of modified Docetaxel, Cisplatin, and fluorouracil (DCF) versus DCF plus growth factor support in patients with metastatic gastric Adenocarcinoma: a study of the US gastric cancer consortium. *J Clin Oncol.* 2015;33:3874–9.
 12. Van Cutsem E, Boni C, Tabernero J, Massuti B, Middleton G, Dane F, et al. Docetaxel plus oxaliplatin with or without fluorouracil or capecitabine in metastatic or locally recurrent gastric cancer: a randomized phase II study. *Ann Oncol.* 2015;26:149–56.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

