

RESEARCH ARTICLE

Open Access

The radiosensitising effect of gemcitabine and its main metabolite dFdU under low oxygen conditions is *in vitro* not dependent on functional HIF-1 protein

An Wouters^{1*}, Bea Pauwels¹, Natalie Burrows^{2,3}, Marc Baay¹, Vanessa Deschoolmeester¹, Trung Nghia Vu⁴, Kris Laukens⁴, Paul Meijnders⁵, Dirk Van Gestel⁵, Kaye J Williams², Danielle Van den Weyngaert⁵, Jan B Vermorken^{1,6}, Patrick Pauwels^{1,7}, Marc Peeters^{1,6} and Filip Lardon¹

Abstract

Background: Regions within solid tumours often experience oxygen deprivation, which is associated with resistance to chemotherapy and irradiation. The aim of this study was to evaluate the radiosensitising effect of gemcitabine and its main metabolite dFdU under normoxia versus hypoxia and to determine whether hypoxia-inducible factor 1 (HIF-1) is involved in the radiosensitising mechanism.

Methods: Stable expression of dominant negative HIF-1 α (dnHIF) in MDA-MB-231 breast cancer cells, that ablated endogenous HIF-1 transcriptional activity, was validated by western blot and functionality was assessed by HIF-1 α activity assay. Cells were exposed to varying oxygen environments and treated with gemcitabine or dFdU for 24 h, followed by irradiation. Clonogenicity was then assessed. Using radiosensitising conditions, cells were collected for cell cycle analysis.

Results: HIF-1 activity was significantly inhibited in cells stably expressing dnHIF. A clear radiosensitising effect under normoxia and hypoxia was observed for both gemcitabine and dFdU. No significant difference in radiobiological parameters between HIF-1 proficient and HIF-1 deficient MDA-MB-231 cells was demonstrated.

Conclusions: For the first time, radiosensitisation by dFdU, the main metabolite of gemcitabine, was demonstrated under low oxygen conditions. No major role for functional HIF-1 protein in radiosensitisation by gemcitabine or dFdU could be shown.

Keywords: Hypoxia, Radiosensitisation, Gemcitabine, dFdU, HIF-1

Background

Regions within solid tumours often experience mild to severe oxygen deprivation (hypoxia) owing to aberrant vascular structure and function. Multiple clinical studies have documented the importance of hypoxia and anoxia (an absence of oxygen) in determining local tumour control in radiotherapy. In addition to the direct role of oxygen in generating radiation-induced DNA damage, the biological effects of hypoxia on tumour cells can also

modulate their response to therapy [1,2]. One major transcription factor involved in the cellular response to reduced oxygen conditions is hypoxia inducible factor 1 (HIF-1). This heterodimeric transcription factor is formed by the association of an oxygen-regulated HIF-1 α subunit with a constitutively expressed HIF-1 β subunit. As HIF-1 modulates many cellular processes, including proliferation, apoptosis, metabolism and the tumour vasculature, it has been reported that HIF-1 has divergent effects on tumour radiosensitivity, which might cause tumours to become more or less radiosensitive [3].

Among the most potent (normoxic) radiosensitisers currently available are antimetabolites. Gemcitabine

* Correspondence: An.Wouters@uantwerpen.be

¹Center for Oncological Research Antwerp, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium

Full list of author information is available at the end of the article

(2',2'-difluorodeoxycytidine, dFdC) is a synthetic pyrimidine nucleoside analogue clinically active against a broad spectrum of solid tumours. Intracellularly, the diphosphate (dFdCDP) and triphosphate (dFdCTP) forms of the drug are responsible for the cytotoxic effects, via inhibition of ribonucleotide reductase and by incorporation into the DNA, leading to chain termination, respectively. In addition to its cytotoxic effect, gemcitabine has potent radiosensitising properties, shown in both preclinical and clinical studies [4]. Current evidence suggests that accumulation in the S phase of the cell cycle, depletion of dATP pools, reduction of apoptotic threshold, inhibition of DNA synthesis and reduction of DNA repair may contribute to, or might even be essential for gemcitabine-mediated radiosensitisation [5].

Following intravenous administration of gemcitabine, the drug rapidly undergoes deamination to its main metabolite, 2',2'-difluorodeoxyuridine (dFdU), resulting in a plasma half-life of gemcitabine of only eight minutes [6]. In contrast, the half-life of dFdU is greater than 14 h, yielding elevated dFdU plasma concentrations for a prolonged period of time (>24 h) at levels known to cause growth inhibition. Importantly, although dFdU has limited cytotoxic activity, it has been demonstrated that it causes a clear concentration- and schedule-dependent radiosensitising effect *in vitro* and potentially contributes to the potent radiosensitising properties of gemcitabine in the clinic [7].

Thus far, few preclinical studies have focused on the outcome of chemoradiation treatments under hypoxia, and on the potential impact of functional HIF-1 on the radiosensitising effect of cytotoxic agents. The molecular basis of hypoxia-mediated chemotherapy and radiotherapy failure indeed has only recently been reported. In these studies, a contribution of HIF-1 to drug resistance has been observed in a wide spectrum of neoplastic cells and many signalling pathways, including PI3K, MAPK, HER2, EGFR and COX2, are reported to induce chemoresistance through HIF-1 activity [8-11].

Concerning gemcitabine, it has recently been observed that this drug radiosensitises both p53 wild type and p53 deficient non-small cell lung cancer cells under hypoxia [12]. Although it was described that gemcitabine did not affect tumour oxygenation or HIF-1 α levels in HCT116 xenografts [13], it has also been reported that gemcitabine inhibited HIF-1 α induction in A549 cells exposed to the hypoxia mimetic agent DFX [14]. In contrast, a more recent study showed gemcitabine-induced activation of HIF-1 α in normoxic pancreatic cancer cells [15]. In order to further elucidate whether or not the HIF-1 transcription factor is involved in the retained radiosensitisation by gemcitabine under low oxygen conditions, in the present study, we evaluated the impact of hypoxia on radiosensitisation by gemcitabine and dFdU in three

isogenic breast adenocarcinoma cell lines differing in HIF-1 status.

Methods

Cell culture

The human tumour cell lines included were MDA-MB-231 (breast adenocarcinoma; wild type (wt) HIF-1) and the sublines MDA-MB-231 dnHIF (dominant-negative HIF-1 α ; HIF-1 activity inhibited) and MDA-MB-231 empty vector control (EV; functional HIF-1). MDA-MB-231 sublines were constructed as described previously [16], resulting in MDA-MB-231 cells stably expressing dnHIF tagged with enhanced green fluorescence protein (eGFP) or eGFP alone (MDA-MB-231 dnHIF and MDA-MB-231 EV, respectively). The dnHIF construct inhibits HIF-1 activity by competing with endogenous HIF-1 α for interaction with HIF-1 β and DNA binding; it is however likely that non-canonical regulation by HIF-1 is not inhibited, since the dnHIF construct is identical to endogenous HIF-1 α except for loss of the oxygen-dependent degradation domains and DNA-binding domains. All cell lines were free from mycoplasma contamination. Cultures were maintained in exponential growth in a humidified 5% CO₂/95% air atmosphere at 37°C (normoxia).

Oxygen conditions

Hypoxia (<0.1% O₂) was achieved in a Bactron IV anaerobic chamber (Shel Lab, Cornelius, USA), as described previously [17]. Hypoxic incubation was initiated after cells had been cultured under normoxia overnight, allowing attachment to culture dishes.

Western blot analysis

Cells were placed under normoxia or hypoxia for 18 h, yielding a robust induction of the expression of HIF-1 α and HIF-1-induced downstream targets. Subsequently, cells were lysed and protocols were used as previously described [18]. In short, cells were lysed in 100 μ l lysis buffer (10 mM Tris (pH 7.4), 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 50 mM NaF, 1 mM sodium orthovanadate, 1% Triton X-100 v/v, 0.5% Nonidet P-40 v/v, 2 mM leupeptin, 0.15 mM aprotinin, 1.46 mM pepstatin, 1 mM phenylmethanesulfonyl fluoride). For western blot analysis, proteins (20 μ g/lane) were resolved on a 7.5% SDS-PAGE gel and electrotransferred onto a polyvinylidene fluoride (PVDF) membrane (Millipore, Schwalbach, Germany) using standard procedures. After blocking with 5% non-fat dry milk w/v in PBS-T (137 mM NaCl, 2.7 mM KCl, 4.3 mM di-sodiumhydrogenphosphate, 1.4 mM potassium-di-hydrogenphosphate, 0.1% Tween20 v/v) overnight at 4°C, the blot was probed with primary antibodies (mouse monoclonal anti-HIF-1 α (BD Transduction Laboratories, Oxford, UK); monoclonal anti-CA9 (clone M75; kindly provided by Dr. Jaromir Pastorek,

Bratislava, Slovakia)). The blot was then reacted with suitable secondary alkaline phosphatase-conjugated antibodies (Dianova, Hamburg, Germany), followed by detection of the protein with CDP-Star chemiluminescent reagent (Applied Biosystems, Foster City, USA). Finally, the blot was stripped and reprobed for β -actin (mouse monoclonal anti- β -actin, Sigma-Aldrich, Dorset, UK) to ensure equal loading and transfer of proteins.

Immunofluorescence

To assess dnHIF-mediated inhibition of CA9, cells were cultured on sterile coverslips under normoxia or hypoxia for 18 h, fixed for 10 min and permeabilised. Cells were incubated with a mouse monoclonal anti-human CA9 antibody (clone M75) for 1 h at 37°C, washed and incubated with a secondary Alexaflour antibody (Life Technologies, Paisley, UK) for 45 min at 37°C. Cells were then washed and nuclei were counterstained with 4',6-diamidino-2-phenylindole (DAPI) and mounted in DAKO fluorescent mounting media (DAKO, Cambridgeshire, UK). Relative localisation of green (eGFP; to identify eGFP in empty vector cells and the dnHIF construct in dnHIF cells), red (for CA9) and blue (for nuclei) fluorescence was analysed with a snapshot wide-field fluorescence microscope and MetaView software.

HIF-1 activity assay

HIF-1 α activity was assessed as described previously [18]. Briefly, cells were transfected with an adenovirus containing trimers of the LDH-A HRE linked to luciferase and were exposed to hypoxia for 18 h. Afterwards, cells were lysed, luciferase activity per μ g protein was calculated and activity of MDA-MB-231 dnHIF cells was normalised to MDA-MB-231 EV control cells.

Vascular endothelial growth factor ELISA

Cells were incubated under normoxia or hypoxia for 18 h, then the media was removed for analysis of secreted VEGF levels as previously described [18]. The concentration of secreted VEGF was determined using a DuoSet Human VEGF ELISA kit (R&D Systems, Abingdon, UK) according to the manufacturer's instructions, and corrected to the amount of protein within the cell cultures from which the medium was taken.

Human hypoxia signalling pathway PCR array

After 24 h incubation under hypoxia, total RNA samples were isolated from 4.10^6 normoxic and hypoxic cells, using RNeasy[®] Mini Kit (Qiagen, Venlo, The Netherlands). The concentration of extracted RNA (A_{260}/A_{280} ratio) and purity (A_{260}/A_{230} ratio) were measured by Nanodrop ND-1000 spectrophotometer (Isogen, Sint-Pieters-Leeuw, Belgium). The quality of isolated RNA was confirmed by capillary electrophoresis with an Agilent 2100 Bioanalyzer

(Agilent Technologies, Amstelveen, The Netherlands). One μ g RNA was reversed transcribed using Reaction-Ready First Strand cDNA Synthesis Kit (Qiagen) in accordance with the manufacturer's instructions. The relative expression of 84 genes related to the hypoxia signalling pathway was assessed by use of the Human Hypoxia Signalling Pathway PCR Array (Qiagen) and the RT² Real-time SYBR Green/Rox PCR Master mix kit (Qiagen).

Chemoradiation clonogenic assay

MDA-MB-231 cells were exposed to normoxia or hypoxia and treated with 0–8 nM gemcitabine or 0–4 μ M dFdU for 24 h immediately before and during irradiation (0–8 Gy, room temperature, XRAD320 irradiator (Precision X-Ray, North Branford, USA)). Treatment schedule and concentrations of gemcitabine and dFdU were chosen based on previous results [12,19]. To irradiate cells under hypoxia, custom-made airtight Perspex shells, pre-incubated in the anaerobic chamber overnight, were used [17]. Immediately following irradiation, hypoxic cells were reoxygenated and all cells were washed with drug-free medium. Unirradiated control cells were handled identically to treated cells. Following an 8-day incubation period, cells were stained with crystal violet and colonies (>50 cells) were counted as described previously [20].

Analysis of the cell cycle distribution

Cells were incubated with 0–8 nM gemcitabine or 0–4 μ M dFdU for 24 h, under normoxia or hypoxia. Cell cycle distribution was monitored according to the Vindelov method, as previously described [12]. Samples were analysed using a FACScan flow cytometer (Becton Dickinson). Histograms of DNA content were analysed using WinMDI software to determine the fractions in each phase of the cell cycle (G_0/G_1 , S and G_2/M).

Statistical analysis

All experiments were performed independently at least three times, and each experiment comprised at least two parallel samples. Results, if not otherwise stated, are presented as mean \pm standard deviation (SD). Statistical differences were evaluated with two-sided two-sample t-tests, one-way ANOVA or two-way ANOVA, using SPSS v16.0 software. Two-way ANOVA was used to study the influence of oxygen tension, HIF functionality and treatment with gemcitabine, dFdU and/or irradiation on the outcome parameter (i.e. cell survival or cell cycle distribution). *Post hoc* comparisons revealed which groups differed significantly from one another. P values less than 0.05 were considered to be statistically significant.

For irradiation experiments, survival rates were calculated as [mean plating efficiency of treated cells/mean plating efficiency of control cells] × 100%. Radiation survival curves were fitted according to the linear-quadratic model using WinNonlin (Pharsight, Mountain View, USA) with survival = exp(-αD-βD²). The following parameters were calculated: ID₅₀ (radiation dose producing a surviving fraction of 50%); SF₂ (surviving fraction at 2 Gy); and mean inactivation dose (MID). The oxygen enhancement ratio (OER) was determined by dividing ID₅₀ under hypoxia by ID₅₀ under normoxia. The dose enhancement factor (DEF) was calculated as ID₅₀ for control, untreated cells divided by ID₅₀ for treated cells.

Possible synergism between gemcitabine or dFdU and radiation was determined by calculation of the combination index (CI) using CalcuSyn software (Biosoft, Cambridge, UK). CI < 1.0, CI = 1.0 and CI > 1.0 indicated synergism, additivity or antagonism, respectively.

To analyse the PCR array data, relative changes in gene expression were calculated using the ΔΔC_t method. Based on the geNorm algorithm, four endogenous control genes were selected for normalisation. Each replicate cycle threshold (C_t) was normalised to the average C_t of the four endogenous controls on a per plate basis and mRNA expression levels were presented as fold changes.

Results

Validation of stable transfection of MDA-MB-231 cells with dnHIF

Exposure to 18 h hypoxia induced expression of HIF-1α and its downstream target carbonic anhydrase 9 (CA9) in MDA-MB-231 wt and EV cells (Figure 1A). In MDA-MB-231 dnHIF cells, HIF-1α expression was detected in hypoxic cells exposed to reduced oxygen levels too. However, due to the presence of the dnHIF protein, CA9 expression remained markedly lower in comparison with the HIF-1α proficient cell lines. As shown in Figure 1B, the dnHIF construct was localised to the nucleus and was expressed independently of oxygen availability. Transfection with the control vector (EV) resulted in eGFP expression that was confined to the cytoplasm. Moreover, the immunofluorescence images clearly showed induction of CA9 in the EV cells under hypoxic conditions, while CA9 staining was absent in hypoxic dnHIF cells.

In order to quantitatively evaluate the impact of dnHIF on HIF-1 function, an adenoviral-based HIF-1α reporter gene assay showed that HIF-activity was significantly inhibited (p < 0.01) in cells expressing the dnHIF protein (Figure 1C). Furthermore, hypoxia-induced VEGF secretion was significantly lower (p < 0.05) in dnHIF versus EV cells (Figure 1D).




Figure 1 dnHIF expression reduces expression of HIF-1 target proteins (CA-9 and VEGF) and HIF-1α activity. **A.** Western blot analysis of HIF-1α, dnHIF and CA9 protein level in MDA-MB-231 wt, EV and dnHIF cells. Cells were exposed to normoxia (N) or hypoxia (H) for 18 h. β-actin detection served as the loading control. **B.** Fluorescent images of MDA-MB-231 EV and dnHIF cells stained with eGFP/dnHIF (green), CA9 (red) and DAPI (blue). Cells were exposed to normoxia or hypoxia for 18 h. **C.** HIF-reporter assay in MDA-MB-231 cells stably-expressing dnHIF or empty vector (EV) control following 18 h hypoxic exposure. Luciferase activity per μg protein was calculated and activity was normalised to EV control cells (**: p < 0.01 in dnHIF vs. EV cells). All results are from at least 3 independent experiments ± SD. **D.** VEGF levels in media taken from dnHIF and EV control cells in normoxia/hypoxia for 18 h. VEGF media levels were normalised to mg protein within the cell cultures from which the media were taken. Hypoxia significantly increased VEGF expression in control MDA-MB-231 EV cells (**: p < 0.01 in hypoxic vs. normoxic cells) but not in dnHIF cells. Most importantly, inhibition of HIF by use of a dominant-negative protein variant (dnHIF) inhibited hypoxia-induced VEGF expression (*: p < 0.05 in dnHIF vs. EV cells). All results are from at least 3 independent experiments ± SD.

Table 1 mRNA expression profiles of hypoxia-related genes in MDA-MB-231 wt, EV and dnHIF cells

Symbol	Description	Fold up- or downregulation under hypoxia versus normoxia		
		MDA-MB-231 wt	MDA-MB-231 EV	MDA-MB-231 dnHIF
ADM	Adrenomedullin	2.18	1.73	-1.00
ANGPTL4	Angiopoetin-like 4	2.72	3.19	2.65
BHLHE40	Basic helix-loop-helix family, member e40	1.56	2.18	2.16
CASP1	Caspase-1	-1.44	-2.03	-1.67
HIF-1 α	Hypoxia inducible factor 1, alpha subunit	-2.61	-3.26	-2.29
HIF-3 α	Hypoxia inducible factor 3, alpha subunit	-2.25	-1.71	-1.68
IPCEF1	Interaction protein for cytohesin exchange factors 1	-3.75	-1.51	4.48
LCT	Lactase	-1.02	-2.39	-1.73
LEP	Leptin	-1.96	-1.04	-4.06
MT3	Metallothionein 3	1.52	1.06	2.01
SLC2A4	Solute carrier family 2 (facilitated glucose transporter), member 4	-2.10	-1.16	-1.38

Only genes with a more than two-fold up- or downregulation under hypoxia versus normoxia are presented. (-) represents downregulation. Cells were exposed to normoxia/hypoxia for 24 h (see Methods for full details). All results are from at least 3 independent experiments.


Human hypoxia signalling pathway PCR array


Out of 84 genes related to the hypoxia signalling pathway, 11 genes showed a more than two-fold up- or down-regulation in mRNA levels between normoxic and hypoxic conditions (Table 1). For HIF-1 α , a decreased mRNA expression level was observed after 24 h hypoxia (Figure 2). Statistical analysis revealed no differential expression profile between MDA-MB-231 wt and EV versus MDA-MB-231 dnHIF cells for the hypoxia-related genes included in the PCR array. A significant difference in expression level under normoxia versus hypoxia was noticed for HIF-1 α and angiopoetin-like 4 in MDA-MB-231 wt

and EV cells ($p < 0.042$); in MDA-MB-231 dnHIF cells, this was only detected for angiopoetin-like 4 ($p = 0.026$). Significance was however lost when p-values were adjusted for multiple comparisons according to the Benjamini-Hochberg procedure.

Cell survival after treatment with gemcitabine or dFdU plus radiation

A concentration-dependent radiosensitising effect was observed for gemcitabine and dFdU (Figure 3, Table 2), yielding a moderately synergistic to synergistic interaction between gemcitabine and irradiation. Radiosensitisation





was similar under normoxia and hypoxia ($p = 0.477$ for gemcitabine, $p = 0.563$ for dFdu). Moreover, the dose enhancement factor was not significantly influenced by the cell line used ($p = 0.736$ for gemcitabine, $p = 0.832$ for dFdu) and a similar degree of radiosensitisation was observed with gemcitabine and dFdu in MDA-MB-231 wt, EV and dnHIF cells. Cell survival was significantly influenced by the concentration of gemcitabine or dFdu, dose of radiation, and oxygen tension ($p \leq 0.001$). While treatment with gemcitabine or dFdu and exposure to hypoxia both had a significant impact on ID_{50} , MID and SF_{2} , no significant difference for these radiobiological parameters was seen between EV versus dnHIF cells.

Cell cycle analysis after 24 h treatment with gemcitabine or dFdu

In contrast to previous findings in other cell lines [17], exposure of MDA-MB-231 cells to hypoxia did not induce a significant increase in the percentage of $G_{0/1}$ cells

($p = 0.213$) (Figure 4). The number of cells in S phase was significantly influenced by the concentration of gemcitabine or dFdu ($p < 0.001$). Gemcitabine and dFdu caused a S phase block in MDA-MB-231 wt, EV and dnHIF cells, both under normoxia and hypoxia (Table 3). As shown on the DNA histograms (Figure 4B), the cell cycle arrest was clearly dependent on the concentration of the drug and shifted from a S phase block to an early S phase block, near the G_1/S border, with increasing concentrations of gemcitabine and dFdu. *Post hoc* analysis revealed no significant difference in the percentage of $G_{0/1}$, S or G_2/M phase cells between MDA-MB-231 EV and MDA-MB-231 dnHIF at any condition tested, suggesting that the cell cycle perturbations were not dependent on functionality of HIF-1 α .

Discussion

The therapeutic implications of oxygen deficiency have been fuelling cancer research for over 100 years. However,

Table 2 Radiobiological parameters for the combination of gemcitabine or dFdU with irradiation under normoxia or hypoxia

Condition	OER	SD	DEF	SD	CI	SD	ID ₅₀	SD	MID	SD	SF ₂	SD	% survival	SD					
MDA-MB-231 wt																			
24 N → RT							2.83	1.03	3.30	1.06	65.61	21.99	100	0					
24 N + 8 nM dFdC → RT			1.70	0.96	0.865	0.175	*	1.70	0.72	††	2.00	0.76	††	37.48	17.47	††	60	27	††
24 N + 2 μM dFdU → RT			2.18	0.57	0.701	0.214	*	1.46	0.22	††	1.62	0.28	††	29.63	9.83	††	41	12	††
24 H → RT	1.43	0.61					3.67	1.03	4.14	0.98	72.96	16.96	93	11					
24 H + 8 nM dFdC → RT			1.59	0.46	0.683	0.243	¶	2.37	1.32	††	2.51	1.02	††	58.92	17.91		60	25	††
24 H + 2 μM dFdU → RT			2.34	0.71	0.687	0.285	¶	1.87	0.59	††	2.34	0.32	†,††	49.39	15.89		44	8	††
MDA-MB-231 EV																			
24 N → RT							2.35	1.29	2.65	1.37	53.68	34.97	100	0					
24 N + 8 nM dFdC → RT			1.70	0.82	0.932	0.273		1.59	0.66		1.61	0.24		28.02	8.46		39	9	††
24 N + 2 μM dFdU → RT			1.74	1.06	0.839	0.231	*	1.80	0.75		2.05	0.72		40.16	26.53		45	23	††
24 H → RT	1.57	0.76					3.41	1.78	4.42	1.81	68.53	32.22	92	36					
24 H + 8 nM dFdC → RT			1.77	0.85	0.381	0.023	¶	2.00	0.47		2.56	1.04		52.56	22.09		40	30	††
24 H + 2 μM dFdU → RT			2.35	1.16	0.711	0.245	*	2.01	0.45		2.59	0.71		48.64	7.71		49		24
MDA-MB-231 dnHIF																			
24 N → RT							3.02	1.42	3.46	1.73	69.79	38.04	100	0					
24 N + 8 nM dFdC → RT			1.50	0.75	0.761	0.182	*	1.82	1.18		2.25	1.16		42.90	26.39		30	15	††
24 N + 2 μM dFdU → RT			2.45	0.67	0.818	0.274	*	1.86	0.99		2.07	1.13		45.32	33.00		61	15	††
24 H → RT	1.43	0.54					3.80	1.40	4.93	1.47	81.34	35.51	97	19					
24 H + 8 nM dFdC → RT			2.04	0.93	0.587	0.280	¶	1.90	0.86	††	1.97	0.69	††	42.62	20.43	††	32	15	††
24 H + 2 μM dFdU → RT			2.08	0.52	0.798	0.311	*	2.54	0.77		2.85	0.83	††	59.16	14.76		63	16	††


dFdC: gemcitabine; RT: irradiation; N: normoxia; H: hypoxia; SD: standard deviation; OER: oxygen enhancement ratio; DEF: dose enhancement factor; CI: combination index, ID₅₀: radiation dose producing a surviving fraction of 50%; MID: mean inactivation dose (MID), SF₂: surviving fraction at 2 Gy; % survival: representing the cytotoxic effect of gemcitabine (dFdC) or dFdU and/or hypoxia alone, 0 Gy.
 †: p < 0.05 vs. corresponding normoxic condition (same treatment, same cell line); ††: p < 0.05 vs. corresponding untreated condition (same pO₂, same cell line).
 *: moderate synergism (0.7 > CI > 0.9); ¶: synergism (CI < 0.7).
 All results are from at least 3 independent experiments ± SD.

detailed studies on the impact of hypoxia on the cytotoxic and/or radiosensitising effects of anticancer drugs are lacking. As the adaptation of tumour cells to hypoxia is primarily mediated by stabilisation of HIF-1, we evaluated the role of functional HIF-1 in the response to chemoradiotherapy. Interestingly, several studies have shown that the presence of HIF-1α is a negative prognostic factor for human breast cancer [21-23]. Animal studies of metastatic breast cancer have demonstrated that lack of HIF-1α in malignant cells significantly reduced tumour progression and metastasis [24]. Moreover, high HIF-1α levels were shown to be predictive of response to epirubicin therapy in patients with breast cancer [25].

In our study, Western blotting showed a consistent upregulation of HIF-1α protein level under hypoxia, whereas the PCR array indicated that HIF-1α was down-regulated on mRNA level under hypoxia. Similarly, exposure of HeLa cells to hypoxia (1% O₂) or the oxygen mimetic CoCl₂ for 2.5 h did not change HIF-1α mRNA levels significantly, while HIF-1α protein levels increased [26]. In patients with colon cancer, high HIF-1 expression

was demonstrated using immunohistochemistry, but no significant difference in HIF-1α mRNA expression between tumour groups and control groups was noticed [27]. One possible explanation for such a discrepancy between mRNA and protein levels is that induction of HIF-1α protein expression is not due to enhanced HIF-1α gene transcription or elevated mRNA stability, but instead results from a longer half-life of the protein due to increased HIF-1α translation and decreased HIF-1α proteolysis [26].

Importantly, due to the above-described transient stabilisation and short half-life of endogenous HIF-1α, HIF targets such as CA9 and the glucose transporter 1 (GLUT-1) have been used to detect hypoxic response in tumour tissues. In breast cancer, abundant expression of CA9 and GLUT-1 was shown to be associated with high-grade cancers and poor prognosis [28,29]. Moreover, CA9 has also been suggested as a predictive marker for response to doxorubicin treatment and adjuvant endocrine therapy in patients with breast cancer [30,31]. In addition, several gene and miRNA expression signatures have been described to be associated with poor prognosis in breast



carcinoma [32]. In this respect, there is a pressing need for better biomarkers of hypoxia (including gene expression profiles, serum proteins, circulating tumour cells or functional imaging) that could be used non-invasively in patients to enable more rigorous testing of its prognostic/predictive value [33].

Concerning the cytotoxic effect of gemcitabine, no significant influence of hypoxia was observed in the breast carcinoma cell lines included in the present study. In addition, for the first time, the effect of gemcitabine's main metabolite dFdU was investigated under reduced oxygen conditions and a similar cytotoxic effect was shown under normoxia and hypoxia. This might be explained by the fact that hypoxia has been shown to have no effect on protein expression of several key enzymes (including dCK and cytidine monophosphate kinase) responsible for metabolism of gemcitabine [34].

Moreover, we noticed that both gemcitabine and dFdU induced a clear S phase block in normoxic and hypoxic cells, independent on HIF-1 functionality. Also, no reduction of cellular uptake and DNA incorporation of gemcitabine

under hypoxia was reported in pancreatic carcinoma and hepatoma-derived cell lines [35].

Other papers however showed that oxygen deficiency did compromise the cytotoxic effect of gemcitabine, suggesting a cell type dependency of this phenomenon. For example, treatment of transitional cell carcinoma cells with gemcitabine was less effective under hypoxia [36]. For pancreatic cancer cells, several studies reported that hypoxia induced resistance to gemcitabine, by altered signalling through PI3K/Akt/NF-κB pathways and partially through MAPK signalling pathway [37], by reducing both inhibition of proliferation and induction of apoptosis by gemcitabine [38], and by decreasing the synthesis of active gemcitabine deoxynucleotides, possibly also through downregulation of dCK [39]. As such, the impact of tumour-associated hypoxia on the cytotoxic effect of gemcitabine is still not completely resolved.

The present report showed no association between radiosensitisation by gemcitabine or dFdU and HIF-1 functionality. Previous work either focused on the relationship

Table 3 Cell cycle distribution after 24 h gemcitabine or dFdU under normoxia or hypoxia

Condition	G _{0/1} phase		S phase		G ₂ /M phase	
	Mean	SD	Mean	SD	Mean	SD
MDA-MB-231 wt						
0 nM dFdC/dFdU (N)	51.36	7.38	24.85	3.39	18.79	4.90
8 nM dFdC (N)	25.76	8.82	67.73	6.35	9.65	1.66
2 μM dFdU (N)	37.25	8.74	44.28	18.19	13.07	8.33
0 nM dFdC/dFdU (H)	48.61	6.04	29.69	7.94	18.41	8.85
8 nM dFdC (H)	41.95	17.94	53.28	17.17	8.81	1.43
2 μM dFdU (H)	49.31	11.21	37.79	12.98	5.46	0.22
MDA-MB-231 EV						
0 nM dFdC/dFdU (N)	44.02	9.86	25.31	5.77	16.31	2.50
8 nM dFdC (N)	43.97	6.20	29.82	7.59	13.53	4.80
2 μM dFdU (N)	39.05	15.49	42.19	19.60	8.72	2.45
0 nM dFdC/dFdU (H)	51.31	11.10	27.28	9.94	10.17	4.03
8 nM dFdC (H)	52.53	13.96	28.11	13.79	8.57	2.37
2 μM dFdU (H)	50.51	14.73	33.15	19.84	6.56	2.31
MDA-MB-231 dnHIF						
0 nM dFdC/dFdU (N)	49.22	8.46	23.60	4.67	16.97	2.19
8 nM dFdC (N)	41.89	18.19	31.65	15.59	13.38	7.75
2 μM dFdU (N)	41.77	18.66	44.09	14.32	7.94	1.98
0 nM dFdC/dFdU (H)	47.65	8.00	28.50	5.61	13.03	3.99
8 nM dFdC (H)	40.86	13.01	34.76	14.29	12.90	8.00
2 μM dFdU (H)	51.22	12.61	34.17	8.23	7.18	1.81

dFdC: gemcitabine; N: normoxia; H: hypoxia.

¶: p < 0.05 vs. corresponding normoxic condition (same treatment, same cell line); *: p < 0.05 vs. untreated control (same pO₂, same cell line).

All results are from at least 3 independent experiments ± SD.

between HIF-1 and the cytotoxic effect of gemcitabine or between HIF-1 and radiosensitivity per se.

Firstly, previous observations have been somewhat controversial regarding HIF-1α expression and the sensitivity to gemcitabine. Suppression of HIF-1α using siRNA resulted in an enhanced efficacy of gemcitabine in the treatment of several pancreatic tumour cell lines [40,41]. Nevertheless, in line with our results, knockdown of HIF-1α has also been reported to have no effect on the sensitivity of pancreatic PANC-1 cells when treated with gemcitabine under hypoxic conditions [35]. In addition, HIF-1α expression levels after platinum/gemcitabine therapy did not correlate with outcome of patients with stage II/III non-small cell lung cancer and HIF-1α expression was not associated with adverse effects or outcome in patients with pancreatic cancer [42]. As such, the therapeutic value of an approach by which gemcitabine is combined with inactivation of HIF-1α signalling by novel strategies remains to be fully elucidated.

Secondly, the effects of HIF-1 blockade on tumour radiosensitivity are complex. Downstream effects of HIF-1 serve to help tumour cells to adapt to hypoxic stress. In doing so, they change the tumour phenotype

in ways that might impact radiosensitivity, some positively and some negatively, but the degree and direction of that influence appears to be dependent on the context. For example, inhibition of HIF-1 activation using siRNA clearly increased radiosensitivity of hypoxic fibrosarcoma cells [43]. Other studies however suggested that HIF-1 would not affect radiosensitivity [44]. From the experimental model used in this study, two conclusions can be drawn. Firstly, no significant difference in radiosensitivity was observed for HIF-1 proficient versus deficient cells, with an OER around 1.50 for all three cell lines. A comparable and relatively low OER of 1.86 ± 0.73 has been reported for MDA-MB-231 by Lagadec *et al.*, who suggested that a negative correlation exists between the OER and increasing malignancy of the breast cancer subtype the cell lines were originally derived from [45]. Secondly, none of the radiobiological parameters (ID₅₀, MID, SF₂) calculated were significantly influenced by HIF-1 functionality after treatment with gemcitabine or dFdU in combination with radiation.

One important limitation of our *in vitro* study is the lack of the microenvironment that would surround tumours *in vivo*. Therefore, further studies using tumour animal

models would certainly be warranted. Only in this way, an in-depth understanding and characterisation of hypoxia in breast cancer and other relevant tumour types can be established, ultimately enabling an enhanced prediction of prognosis, optimisation of (gemcitabine and/or radiation) treatment and information on whether and how to target tumour hypoxia.

Conclusions

Taking into account our previous work in lung cancer cell lines [12], this study showed that the retained radiosensitising effect of gemcitabine under hypoxia was not tumour tissue specific and could be observed in MDA-MB-231 breast cancer cells. For the first time, radiosensitisation by dFdU, the main metabolite of gemcitabine, was investigated and demonstrated under low oxygen conditions. As dFdU has a prolonged half-life, the sustained presence of dFdU in the blood might induce radiosensitisation despite the short half-life of the parent drug gemcitabine. This might be highly relevant, especially considering delivery of the drug to hypoxic tumour regions. As HIF-1 proficient and HIF-1 deficient cells were equally radiosensitised, no major role for functional HIF-1 protein in radiosensitisation by gemcitabine or dFdU could be demonstrated.

Competing interests

Actual or potential conflicts of interest do not exist.

Authors' contributions

AW carried out the cell culture experiments and drafted the manuscript. BP carried out the PCR array and helped to draft the manuscript. NB and KJW carried out the transfections and the molecular biological studies. MB and VD participated in the design of the study and helped to draft the manuscript. TNV and KL performed the statistical analysis. PM, DVG and DVDW participated in the radiation experiments. JBV, PP, MP and FL participated in the study design and coordination. All authors read and approved the final manuscript.

Acknowledgements

AW is funded by Research Foundation Flanders (FWO-Vlaanderen, Belgium) as postdoctoral fellow. Gemcitabine and dFdU were kindly provided by Eli Lilly Company.

Author details

¹Center for Oncological Research Antwerp, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium. ²Hypoxia and Therapeutics Group, University of Manchester, Oxford Road, Manchester M13 9PT, UK. ³Cambridge Institute for Medical Research, University of Cambridge, 4.17 Wellcome Trust/MRC Building, Addenbrooke's Hospital, Hills Road, Cambridge CB2 0XY, UK. ⁴Biomedical Informatics Research Center Antwerp (Biomina), University of Antwerp, Middelheimlaan 1, 2020 Antwerpen, Belgium. ⁵Department of Radiotherapy, University Radiotherapy Antwerp (URA), Lindendreef 1, 2020 Antwerp, Belgium. ⁶Department of Oncology, Antwerp University Hospital, Wilrijkstraat 10, 2650 Edegem, Belgium. ⁷Department of Pathology, Antwerp University Hospital, Wilrijkstraat 10, 2650 Edegem, Belgium.

Received: 8 April 2014 Accepted: 5 August 2014

Published: 16 August 2014

References

1. Bertout JA, Patel SA, Simon MC: **The impact of O₂ availability on human cancer.** *Nat Rev Cancer* 2008, **8**:967–975.

- Wouters A, Pauwels B, Lardon F, Vermorken JB: **Review: implications of in vitro research on the effect of radiotherapy and chemotherapy under hypoxic conditions.** *Oncologist* 2007, **12**:690–712.
- Moeller B, Dewhirst M: **HIF-1 and tumour radiosensitivity.** *Br J Cancer* 2006, **95**:1–5.
- Pauwels B, Korst A, Lardon F, Vermorken J: **Combined modality therapy of gemcitabine and radiation.** *Oncologist* 2005, **10**:34–51.
- Pauwels B, Vermorken JB, Wouters A, Ides J, Van LS, Lambrechts HA, Pattyn GG, Vermeulen K, Meijnders P, Lardon F: **The role of apoptotic cell death in the radiosensitising effect of gemcitabine.** *Br J Cancer* 2009, **101**:628–636.
- Abbruzzese JL, Grunewald R, Weeks EA, Gravel D, Adams T, Nowak B, Mineishi S, Tarassoff P, Satterlee W, Raber MN, Plunkett W: **A phase I clinical, plasma, and cellular pharmacology study of gemcitabine.** *J Clin Oncol* 1991, **9**:491–498.
- Specenier P, Guetens G, Dyck J, de Boeck G, Weyler J, Van den Weyngaert D, Aelbrecht K, Vermorken JB: **Difluorodeoxyuridine plasma concentrations after low-dose gemcitabine during chemoradiation in head and neck cancer patients.** *Cancer Chemother Pharmacol* 2011, **68**:185–191.
- Sasabe E, Zhou X, Li D, Oku N, Yamamoto T, Osaki T: **The involvement of hypoxia-inducible factor-1alpha in the susceptibility to gamma-rays and chemotherapeutic drugs of oral squamous cell carcinoma cells.** *Int J Cancer* 2007, **120**:268–277.
- Hao J, Song X, Song B, Liu Y, Wei L, Wang X, Yu J: **Effects of lentivirus-mediated HIF-1alpha knockdown on hypoxia-related cisplatin resistance and their dependence on p53 status in fibrosarcoma cells.** *Cancer Gene Ther* 2008, **15**:449–455.
- Rohwer N, Cramer T: **Hypoxia-mediated drug resistance: novel insights on the functional interaction of HIFs and cell death pathways.** *Drug Resist Updat* 2011, **14**:191–201.
- Luwor RB, Lu Y, Li X, Mendelsohn J, Fan Z: **The anti-epidermal growth factor receptor monoclonal antibody cetuximab/C225 reduces hypoxia-inducible factor-1 alpha, leading to transcriptional inhibition of vascular endothelial growth factor expression.** *Oncogene* 2005, **24**:4433–4441.
- Wouters A, Pauwels B, Lambrechts HA, Pattyn GG, Ides J, Baay M, Meijnders P, Peeters M, Vermorken JB, Lardon F: **Retention of the in vitro radiosensitizing potential of gemcitabine under anoxic conditions, in p53 wild-type and p53-deficient non-small-cell lung carcinoma cells.** *Int J Radiat Oncol Biol Phys* 2011, **80**:558–566.
- Huxham LA, Kyle AH, Baker JH, Nyklichuk LK, Minchinton AI: **Microregional effects of gemcitabine in HCT-116 xenografts.** *Cancer Res* 2004, **64**:6537–6541.
- Ikeda R, Vermeulen LC, Jiang Z, Lau E, Kolesar JM: **Gemcitabine and paclitaxel suppress the production of vascular endothelial growth factor induced by deferoxamine in human non-small cell lung cancer A549 cells.** *Exp Ther Med* 2010, **1**:853–857.
- Arora S, Bhardwaj A, Singh S, Srivastava SK, McClellan S, Nirodi CS, Piazza GA, Grizzle WE, Owen LB, Singh AP: **An undesired effect of chemotherapy: gemcitabine promotes pancreatic cancer cell invasiveness through reactive oxygen species-dependent, nuclear factor kappaB- and hypoxia-inducible factor 1alpha-mediated up-regulation of CXCR4.** *J Biol Chem* 2013, **288**:21197–21207.
- Roberts DL, Williams KJ, Cowen RL, Barathova M, Eustace AJ, Brittain-Dissont S, Tilby MJ, Pearson DG, Ottley CJ, Stratford IJ, Dive C: **Contribution of HIF-1 and drug penetrance to oxaliplatin resistance in hypoxic colorectal cancer cells.** *Br J Cancer* 2009, **101**:1290–1297.
- Wouters A, Pauwels B, Lambrechts HA, Pattyn GG, Ides J, Baay M, Meijnders P, Dewilde S, Vermorken JB, Lardon F: **Chemoradiation interactions under reduced oxygen conditions: Cellular characteristics of an in vitro model.** *Cancer Lett* 2009, **286**:180–188.
- Burrows N, Babur M, Resch J, Ridsdale S, Mejin M, Rowling EJ, Brabant G, Williams KJ: **GDC-0941 inhibits metastatic characteristics of thyroid carcinomas by targeting both the phosphoinositide-3 kinase (PI3K) and hypoxia-inducible factor-1alpha (HIF-1alpha) pathways.** *J Clin Endocrinol Metab* 2011, **96**:E1934–E1943.
- Pauwels B, Korst AE, Lambrechts HA, Pattyn GG, De Pooter CM, Lardon F, Vermorken JB: **The radiosensitising effect of difluorodeoxyuridine, a metabolite of gemcitabine, in vitro.** *Cancer Chemother Pharmacol* 2006, **58**:219–228.
- Wouters A, Pauwels B, Lambrechts HA, Pattyn GG, Ides J, Baay M, Meijnders P, Lardon F, Vermorken JB: **Counting clonogenic assays from normoxic**

- and anoxic irradiation experiments manually or by using densitometric software. *Phys Med Biol* 2010, **55**:N167–N178.
21. Bos R, van der Groep P, Greijer AE, Shvarts A, Meijer S, Pinedo HM, Semenza GL, van Diest PJ, van der Wall E: **Levels of hypoxia-inducible factor-1alpha independently predict prognosis in patients with lymph node negative breast carcinoma.** *Cancer* 2003, **97**:1573–1581.
 22. Dales JP, Garcia S, Meunier-Carpentier S, Andrac-Meyer L, Haddad O, Lavaut MN, Allasia C, Bonnier P, Charpin C: **Overexpression of hypoxia-inducible factor HIF-1alpha predicts early relapse in breast cancer: retrospective study in a series of 745 patients.** *Int J Cancer* 2005, **116**:734–739.
 23. Schindl M, Schoppmann SF, Samonigg H, Hausmaninger H, Kwasny W, Gnatt M, Jakesz R, Kubista E, Birner P, Oberhuber G, Austrian B, Colorectal Cancer Study G: **Overexpression of hypoxia-inducible factor 1alpha is associated with an unfavorable prognosis in lymph node-positive breast cancer.** *Clin Cancer Res* 2002, **8**:1831–1837.
 24. Liao D, Corle C, Seagroves TN, Johnson RS: **Hypoxia-inducible factor-1alpha is a key regulator of metastasis in a transgenic model of cancer initiation and progression.** *Cancer Res* 2007, **67**:563–572.
 25. Generali D, Berruti A, Brizzi MP, Campo L, Bonardi S, Wigfield S, Bersiga A, Allevi G, Milani M, Aguggini S, Gandolfi V, Dogliotti L, Bottini A, Harris AL, Fox SB: **Hypoxia-inducible factor-1alpha expression predicts a poor response to primary chemoendocrine therapy and disease-free survival in primary human breast cancer.** *Clin Cancer Res* 2006, **12**:4562–4568.
 26. Galban S, Kuwano Y, Pullmann R Jr, Martindale JL, Kim HH, Lal A, Abdelmohsen K, Yang X, Dang Y, Liu JO, Lewis SM, Holcik M, Gorospe M: **RNA-binding proteins HuR and PTB promote the translation of hypoxia-inducible factor 1alpha.** *Mol Cell Biol* 2008, **28**:93–107.
 27. Wu Y, Jin M, Xu H, Shimin Z, He S, Wang L, Zhang Y: **Clinicopathologic significance of HIF-1alpha, CXCR4, and VEGF expression in colon cancer.** *Clin Dev Immunol* 2010, **2010**: doi:10.1155/2010/537531.
 28. Beketic-Oreskovic L, Ozretic P, Rabbani ZN, Jackson IL, Sarcevic B, Levanat S, Maric P, Babic I, Vujaskovic Z: **Prognostic significance of carbonic anhydrase IX (CA-IX), endoglin (CD105) and 8-hydroxy-2'-deoxyguanosine (8-OHdG) in breast cancer patients.** *Pathol Oncol Res* 2011, **17**:593–603.
 29. Pinheiro C, Sousa B, Albergaria A, Paredes J, Duflou R, Vieira D, Schmitt F, Baltazar F: **GLUT1 and CAIX expression profiles in breast cancer correlate with adverse prognostic factors and MCT1 overexpression.** *Histol Histopathol* 2011, **26**:1279–1286.
 30. Span PN, Bussink J, Manders P, Beex LV, Sweep CG: **Carbonic anhydrase-9 expression levels and prognosis in human breast cancer: association with treatment outcome.** *Br J Cancer* 2003, **89**:271–276.
 31. Betof AS, Rabbani ZN, Hardee ME, Kim SJ, Broadwater G, Bentley RC, Snyder SA, Vujaskovic Z, Oosterwijk E, Harris LN, Horton JK, Dewhirst MW, Blackwell KL: **Carbonic anhydrase IX is a predictive marker of doxorubicin resistance in early-stage breast cancer independent of HER2 and TOP2A amplification.** *Br J Cancer* 2012, **106**:916–922.
 32. Favaro E, Lord S, Harris AL, Buffa FM: **Gene expression and hypoxia in breast cancer.** *Genome Med* 2011, **3**:55.
 33. Eccles SA, Aboagye EO, Ali S, Anderson AS, Armes J, Berditchevski F, Blaydes JP, Brennan K, Brown NJ, Bryant HE, Bundred NJ, Burchell JM, Campbell AM, Carroll JS, Clarke RB, Coles CE, Cook GJ, Cox A, Curtin NJ, Dekker LV, Silva Idos S, Duffy SW, Easton DF, Eccles DM, Edwards DR, Edwards J, Evans D, Fenlon DF, Flanagan JM, Foster C, et al: **Critical research gaps and translational priorities for the successful prevention and treatment of breast cancer.** *Breast Cancer Res* 2013, **15**:R92.
 34. Lam W, Bussom S, Cheng YC: **Effect of hypoxia on the expression of phosphoglycerate kinase and antitumor activity of troxacitabine and gemcitabine in non-small cell lung carcinoma.** *Mol Cancer Ther* 2009, **8**:415–423.
 35. Onozuka H, Tsuchihara K, Esumi H: **Hypoglycemic/hypoxic condition in vitro mimicking the tumor microenvironment markedly reduced the efficacy of anticancer drugs.** *Cancer Sci* 2011, **102**:975–982.
 36. Fechner G, Dederichs F, Schmidt D, Muller S, Vaupel P, Albers P: **Hyperoxia-induced improvement of the in vitro response to gemcitabine in transitional cell carcinoma.** *Anticancer Res* 2005, **25**:3413–3418.
 37. Yokoi K, Fidler IJ: **Hypoxia increases resistance of human pancreatic cancer cells to apoptosis induced by gemcitabine.** *Clin Cancer Res* 2004, **10**:2299–2306.
 38. Zhang X, Galardi E, Duquette M, Lawler J, Parangi S: **Antiangiogenic treatment with three thrombospondin-1 type 1 repeats versus gemcitabine in an orthotopic human pancreatic cancer model.** *Clin Cancer Res* 2005, **11**:5622–5630.
 39. Maftouh M, Avan A, Sciarillo R, Granchi C, Leon LG, Rani R, Funel N, Smid K, Honeywell R, Boggi U, Minutolo F, Peters GJ, Giovannetti E: **Synergistic interaction of novel lactate dehydrogenase inhibitors with gemcitabine against pancreatic cancer cells in hypoxia.** *Br J Cancer* 2014, **110**:172–182.
 40. Yang L, Kang WK: **The effect of HIF-1alpha siRNA on growth and chemosensitivity of MIA-paca cell line.** *Yonsei Med J* 2008, **49**:295–300.
 41. Cheng ZX, Wang DW, Liu T, Liu WX, Xia WB, Xu J, Zhang YH, Qu YK, Guo LQ, Ding L, Hou J, Zhong ZH: **Effects of the HIF-1alpha and NF-kappaB loop on epithelialmesenchymal transition and chemoresistance induced by hypoxia in pancreatic cancer cells.** *Oncol Rep* 2014, **31**:1891–1898.
 42. Kasuya K, Tsuchida A, Nagakawa Y, Suzuki M, Abe Y, Itoi T, Serizawa H, Nagao T, Shimazu M, Aoki T: **Hypoxia-inducible factor-1alpha expression and gemcitabine chemotherapy for pancreatic cancer.** *Oncol Rep* 2011, **26**:1399–1406.
 43. Staab A, Fleischer M, Loeffler J, Said HM, Katzer A, Plathow C, Einsele H, Flentje M, Vordermark D: **Small interfering RNA targeting HIF-1alpha reduces hypoxia-dependent transcription and chemoresistance hypoxic HT 1080 human fibrosarcoma cells in vitro.** *Strahlenther Onkol* 2011, **187**:252–259.
 44. Arvold ND, Guha N, Wang D, Matli M, Deen D, Warren R, Haas Kogan D: **Hypoxia-induced radioresistance is independent of hypoxia-inducible factor-1A in vitro.** *Int J Radiat Oncol Biol Phys* 2005, **62**:207–212.
 45. Lagadec C, Dekmezian C, Bauche L, Pajonk F: **Oxygen levels do not determine radiation survival of breast cancer stem cells.** *PLoS One* 2012, **7**:e34545.

doi:10.1186/1471-2407-14-594

Cite this article as: Wouters et al.: The radiosensitising effect of gemcitabine and its main metabolite dFdU under low oxygen conditions is *in vitro* not dependent on functional HIF-1 protein. *BMC Cancer* 2014 **14**:594.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

