

CORRECTION

Open Access



Correction to: Cervical and breast cancer screening uptake among women with serious mental illness: a data linkage study

Charlotte Woodhead^{1†}, Ruth Cunningham^{2†}, Mark Ashworth³, Elizabeth Barley⁴, Robert J. Stewart^{1*} and Max J. Henderson¹

Correction to: BMC Cancer (2016) 16:819

<https://doi.org/10.1186/s12885-016-2842-8>

Following publication of the original article [1], the authors notified us of an error in the reported percentages in Table 3.

Author details

¹Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK. ²Department of Public Health, University of Otago, Wellington, New Zealand. ³Department of Primary Care and Public Health Sciences, King's College London, London, UK. ⁴Faculty of Nursing and Midwifery, King's, College London, London, UK.

Received: 7 February 2019 Accepted: 7 February 2019

Published online: 15 February 2019

Reference

1. Woodhead C, et al. Cervical and breast cancer screening uptake among women with serious mental illness: a data linkage study. *BMC Cancer*. 2016; 16:819. <https://doi.org/10.1186/s12885-016-2842-8>.

* Correspondence: robert.stewart@kcl.ac.uk

†Charlotte Woodhead and Ruth Cunningham contributed equally to this work.

¹Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK

Full list of author information is available at the end of the article



Table 3 Associations between serious mental illness (SMI) status and recent receipt of breast and/or cervical screening overall and by SMI characteristic sub-group

| | Mammography eligible population (N=26,010) | | | | Cervical smear eligible population (N=107,947) | | | |
|--|--|-----------------------------------|--|--|--|-----------------------------------|--|--|
| | Recorded mammography in last 3 years n (%) | Unadjusted OR (95% CI) | Adjusted for socio-demographics OR ^a (95% CI) | Additionally adjusted for consultation rate OR ^b (95% CI) | Recorded cervical smear in last 3/5 yrs n (%) | Unadjusted OR (95% CI) | Adjusted for socio-demographics OR ^a (95% CI) | Additionally adjusted for consultation rate OR ^b (95% CI) |
| Non-SMI | 14205 (56.0) | 1.00 | 1.00 | 1.00 | 67823 (63.7) | 1.00 | 1.00 | 1.00 |
| SMI overall | 305 (48.8) | 0.72 (0.61 - 0.86) ^{***} | 0.69 (0.57 - 0.84) ^{***} | 0.60 (0.49 - 0.73) ^{***} | 848 (60.9) | 1.16 (0.99 - 1.35) | 0.72 (0.60 - 0.85) ^{***} | 0.35 (0.29 - 0.42) ^{***} |
| SMI by diagnosis | | | | | | | | |
| Schizophrenia | 136 (47.9) | 0.64 (0.50 - 0.82) ^{***} | 0.59 (0.45 - 0.78) ^{***} | 0.52 (0.40 - 0.69) ^{***} | 270 (57.3) | 0.95 (0.74 - 1.22) | 0.48 (0.36 - 0.63) ^{***} | 0.24 (0.18 - 0.32) ^{***} |
| Bipolar affective disorder | 67 (50.4) | 0.85 (0.58 - 1.26) | 0.89 (0.59 - 1.35) | 0.72 (0.47 - 1.10) | 231 (67.9) | 1.94 (1.35 - 2.78) ^{***} | 1.23 (0.84 - 1.80) | 0.50 (0.33 - 0.74) ^{**} |
| Other non-organic psychoses | 34 (41.5) | 0.54 (0.33 - 0.87) [*] | 0.53 (0.31 - 0.90) [*] | 0.47 (0.27 - 0.80) ^{**} | 153 (56.7) | 0.82 (0.60 - 1.13) | 0.57 (0.40 - 0.80) ^{**} | 0.33 (0.22 - 0.47) ^{***} |
| Depot injectable | | | | | | | | |
| No | 191 (53.4) | 0.94 (0.74 - 1.20) | 0.97 (0.75 - 1.26) | 0.83 (0.64 - 1.09) | 524 (63.1) | 0.32 (1.08 - 1.62) ^{**} | 0.82 (0.65 - 1.02) | 0.36 (0.29 - 0.46) ^{***} |
| Yes | 80 (42.6) | 0.50 (0.37 - 0.68) ^{***} | 0.46 (0.33 - 0.64) ^{***} | 0.39 (0.27 - 0.54) ^{***} | 199 (58.5) | 0.92 (0.69 - 1.22) | 0.48 (0.35 - 0.66) ^{***} | 0.26 (0.18 - 0.36) ^{***} |
| Any indicator of severity ¹ | | | | | | | | |
| No | 168 (53.7) | 1.00 (0.77 - 1.29) | 0.91 (0.69 - 1.21) | 0.79 (0.59 - 1.06) | 411 (62.7) | 1.32 (1.05 - 1.66) [*] | 0.82 (0.63 - 1.06) | 0.40 (0.31 - 0.53) ^{***} |
| Yes | 137 (43.9) | 0.54 (0.43 - 0.69) ^{***} | 0.54 (0.42 - 0.70) ^{***} | 0.46 (0.36 - 0.61) ^{***} | 437 (59.2) | 1.04 (0.85 - 1.27) | 0.65 (0.52 - 0.81) ^{***} | 0.31 (0.25 - 0.40) ^{***} |
| Any indicator of risk ² | | | | | | | | |
| No | 205 (51.4) | 0.84 (0.67 - 1.05) | 0.81 (0.63 - 1.03) | 0.70 (0.55 - 0.90) ^{**} | 535 (62.9) | 1.31 (1.07 - 1.60) ^{**} | 0.79 (0.63 - 1.00) [*] | 0.38 (0.30 - 0.49) ^{***} |
| Yes | 100 (44.2) | 0.56 (0.42 - 0.74) ^{***} | 0.53 (0.39 - 0.73) ^{***} | 0.46 (0.34 - 0.63) ^{***} | 313 (57.6) | 0.97 (0.77 - 1.23) | 0.62 (0.48 - 0.80) ^{***} | 0.31 (0.24 - 0.41) ^{***} |

Eligible population includes non-linked non-SMI and linked SMI group

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ ¹ Includes any of: ever had an inpatient stay, any record of being treated under the Mental Health Act, any record of difficulty managing their physical health, or any record of an Assertive Outreach/Crisis/A&E episode² Includes any of: recorded history of violence, recorded history of non-compliance, and any record of a forensic history^a Adjusted for age (continuous), ethnicity, and borough-level deprivation^b Additionally adjusted for mean annual number of primary consultations