

CORRECTION

Open Access



Correction: CYP1B1 promotes tumorigenesis via altered expression of CDC20 and DAPK1 genes in renal cell carcinoma

Yozo Mitsui^{1,2*†}, Inik Chang^{3,4†}, Shinichiro Fukuhara⁴, Miho Hiraki¹, Naoko Arichi¹, Hiroaki Yasumoto¹, Hiroshi Hirata², Soichiro Yamamura², Varahram Shahryari², Guoren Deng², Darryn K. Wong², Shahana Majid², Hiroaki Shiina¹, Rajvir Dahiya² and Yuichiro Tanaka^{2*}

Correction: *BMC Cancer* 15, 942 (2015)
<https://doi.org/10.1186/s12885-015-1951-0>

Published online: 25 July 2022

Following publication of the original article [1], the authors identified an error in Fig. 2C.

In Fig. 2C, the wrong image representing wound healing in CYP1B1-siRNA#1-treated Caki-1 cells at 0 hours was inadvertently applied. Although the wrong image was mistakenly utilized, results are essentially the same and does not change the outcome or conclusions of the article. The correct Fig. 2C is given in this correction article.

Reference

1. Mitsui Y, Chang I, Fukuhara S, et al. CYP1B1 promotes tumorigenesis via altered expression of CDC20 and DAPK1 genes in renal cell carcinoma. *BMC Cancer*. 2015;15:942. <https://doi.org/10.1186/s12885-015-1951-0>.

Author details

¹Department of Urology Shimane University Faculty of Medicine, 89-1 Enya-cho, Izumo 693-8501, Japan. ²Department of Urology, San Francisco Veterans Affairs Medical Center and University of California San Francisco, Bldg 42 Rm 109, San Francisco, CA 94121, USA. ³Department of Oral Biology, Yonsei University College of Dentistry, Seoul, South Korea. ⁴Department of Urology, Osaka University Graduate School of Medicine, Suita 565-0871, Japan.

The original article can be found online at <https://doi.org/10.1186/s12885-015-1951-0>.

[†]Yozo Mitsui and Inik Chang contributed equally to this work.

*Correspondence: mitsui@med.shimane-u.ac.jp; Yuichiro.tanaka@ucsf.edu

¹ Department of Urology Shimane University Faculty of Medicine, 89-1 Enya-cho, Izumo 693-8501, Japan

² Department of Urology, San Francisco Veterans Affairs Medical Center and University of California San Francisco, Bldg 42 Rm 109, San Francisco, CA 94121, USA

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

C




Fig. 2 c Representative images of wound healing assay. After siRNA transfection for 48 h, a wound was formed by scraping and closure of wound measured after 24 h. Attenuation of CYP1B1 significantly inhibited cell migration. **, $P < 0.01$. ***, $P < 0.001$